

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 January 2001 (18.01.2001)

PCT

(10) International Publication Number
WO 01/04264 A2

- (51) International Patent Classification⁷: C12N Michael, G. [CA/US]; 1050 Borregas Avenue, #80, Sunnyvale, CA 94089 (US).
- (21) International Application Number: PCT/US00/17887
- (22) International Filing Date: 28 June 2000 (28.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/349,015 7 July 1999 (07.07.1999) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 09/349,015 (CIP)
Filed on 7 July 1999 (07.07.1999)
- (71) Applicant (for all designated States except US): INCYTE GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): JONES, Karen, Ann [GB/GB]; Cregneish, Elmdon, Saffron Walden, Essex CB11 4LT (GB). VOLKMUTH, Wayne [US/US]; 3620 El Encanto, Calabasas, CA 91302 (US). WALKER,
- (74) Agents: MURRY, Lynn, E. et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ATHEROSCLEROSIS-ASSOCIATED GENES

(57) Abstract: The invention provides novel atherosclerosis-associated polynucleotides and polypeptides encoded by those genes. The invention also provides expression vectors, host cells, and antibodies. The invention also provides methods for screening or purifying ligands and diagnosing or treating atherosclerosis.

WO 01/04264 A2

ATHEROSCLEROSIS-ASSOCIATED GENES

TECHNICAL FIELD

The invention relates to 34 atherosclerosis-associated polynucleotides identified by their co-
5 expression with known atherosclerosis genes and their corresponding gene products. The invention also relates to the use of these biomolecules in diagnosis, prognosis, prevention, treatment, and evaluation of therapies for diseases associated with atherosclerosis.

BACKGROUND ART

Atherosclerosis is a disorder characterized by cellular changes in the arterial intima and the
10 formation of arterial plaques containing intra- and extracellular deposits of lipids. The resultant thickening of artery walls and the narrowing of the arterial lumen is the underlying pathologic condition in most cases of coronary artery disease, aortic aneurysm, peripheral vascular disease, and stroke. A cascade of molecules is involved in the cellular morphogenesis, proliferation, and cellular migration which results in an atherosclerotic lesion (Libby *et al.* (1997) *Int J Cardiol* 62:23-29).

15 A healthy artery consists of three layers. The vascular intima, lined by a monolayer of endothelial cells in contact with the blood, contains smooth muscle cells in extracellular matrix. An internal elastic lamina forms the border between the intima and the tunica media. The media contains layers of smooth muscle cells surrounded by a collagen and elastin-rich extracellular matrix. An external elastic lamina forms the border between the media and the adventitia. The adventitia contains
20 nerves and some mast cells and is the origin of the vasa vasorum which supplies blood to the outer layers of the tunica media.

Initiation of an atherosclerotic lesion often occurs following vascular endothelial cell injury as a result of hypertension, diabetes mellitus, hyperlipidemia, fluctuating shear stress, smoking, or transplant rejection. The injury results in the local release of nitric oxide and superoxide anions which
25 react to form cytotoxic peroxynitrite radicals, causing injury to the endothelium and myocytes of the intima. This cellular injury leads to the expression of a variety of molecules that produce local and systemic effects. The initial cellular response to injury includes the release of mediators of inflammation such as cytokines, complement components, prostaglandins, and downstream transcription factors. These molecules promote monocyte infiltration of the vascular intima and lead to
30 the upregulation of adhesion molecules which encourage attachment of the monocytes to the damaged endothelial cells. Additionally, components of the extracellular matrix including collagens, fibrinogens, and matrix Gla protein are induced and provide sites for monocyte attachment. Annexins, plasminogen activator inhibitor 1, and nitric oxide synthases are triggered to counteract these effects.

Monocytes that infiltrate the lesion accumulate modified low density lipoprotein lipid through
35 scavenger receptors such as CD36 and macrophage scavenger receptor type I. The abundance of modified lipids is a factor in atherogenesis and is influenced by modifying enzymes such as lipoprotein

lipase, carboxyl ester lipase, serum amyloid P component, LDL-receptor related protein, microsomal triglyceride transfer protein, and serum esterases such as paraoxonase. Lipid metabolism is governed by cholesterol biosynthesis enzymes such as 3-hydroxy-3-methylglutaryl coenzyme A synthase, and products of the apolipoprotein genes. Modified lipid stabilization and accumulation is aided by
5 perilipin and alpha-2-macroglobulin.

As monocytes accumulate in the lesion, they can rupture and release free cholesterol, cytokines, and procoagulants into the surrounding environment. This process leads to the development of a plaque which consists of a mass of lipid-engorged monocytes and a lipid-rich necrotic core covered by a fibrous cap. The gradual progression of plaque growth is punctuated by thrombus formation which
10 leads to clinical symptoms such as unstable angina, myocardial infarction, or stroke. Thrombus formation is initiated by episodic plaque rupture which exposes flowing blood to tissue factors, which induce coagulation, and collagen, which activates platelets. After initiation of the atherosclerotic lesion, enzymes that degrade extracellular matrix components such as matrix metalloproteinases and cathepsin K are up-regulated, and their inhibitors are down-regulated. This results in destabilization of
15 the atherosclerotic lesion and subsequent complications including myocardial infarction, angina, and stroke. Further arterial occlusion and infiltration increase with the expression of coagulation factors and down-regulation of their inhibitors, antithrombin III, and lipoprotein-associated coagulation inhibitor.

Smooth muscle cells build up in the arterial media and constitute one of the principal cell types
20 in atherosclerotic and restenotic lesions. They show a high degree of plasticity and are able to shift between a differentiated, contractile phenotype and a less differentiated, synthetic phenotype. This modulation occurs as a response to factors secreted from cells at the site of vascular injury and results in structural reorganization with a loss of myofilaments and the formation of an extensive endoplasmic reticulum and a large Golgi complex. Genes encoding secreted protein, acidic and rich in cysteine
25 (SPARC) and endothelin-1 contribute to these changes. At the same time, the expression of cytoskeletal proteins such as calponin, myosin, desmin, and other gene products in the cells is altered. As a result, the smooth muscle cells lose their contractility and become able to migrate from the media to the intima, to proliferate, and to secrete extracellular matrix components which contribute to arterial intimal thickening.

30 The initiation and progression of atherosclerotic lesion development requires the interplay of various molecular pathways. Many genes that participate in these processes are known, and some of them have been shown to have a direct role in atherosclerosis pathogenesis by animal model experiments, *in vitro* assays, and epidemiological studies (Krettek *et al.* (1997) *Arterioscler Thromb Vasc Biol* 17:2897-2903; Fisher *et al.* (1997) *Atherosclerosis* 135:145-159; Shih *et al.* (1998) *Circulation* 95:2684-2693; and Bocan *et al.* (1998) *Atherosclerosis* 139:21-30).

The present invention satisfies a need in the art by providing new compositions that are useful for diagnosis, prognosis, treatment, prevention, and evaluation of therapies for diseases associated with atherosclerosis. We have implemented a method for analyzing gene expression patterns and have identified 34 atherosclerosis-associated polynucleotides through their co-expression with 66 known atherosclerosis-associated genes.

SUMMARY OF THE INVENTION

The invention provides for a substantially purified polynucleotide comprising a gene that is coexpressed with one or more known atherosclerosis-associated genes in a biological sample. Known atherosclerosis-associated genes include and encode human 22kDa smooth muscle protein, calponin, desmin, smooth muscle myosin heavy chain, alpha tropomyosin, human tissue inhibitor of metalloproteinase 3, human tissue inhibitor of metalloproteinase-2, human tissue inhibitor of metalloproteinase-4, pro alpha 1(I) collagen, collagen alpha-2 type I, collagen alpha-6 type I, procollagen alpha 2(V), collagen VI alpha-2, type VI collagen alpha3, pro-alpha-1 type 3 collagen, pro-alpha-1 (V) collagen, collagenase type IV/ matrix metalloproteinase 9/gelatinase B, matrix Gla protein, cathepsin K, fibrinogen beta chain gene, fibrinogen gamma chain gene, pre-pro-von Willebrand factor, coagulation factor II/ prothrombin, coagulation factor XII, coagulation factor VII, platelet endothelial cell adhesion molecule, lipoprotein-associated coagulation inhibitor, antithrombin III variant, plasminogen activator inhibitor-1, lipoprotein lipase, alpha-2-macroglobulin, apolipoprotein AI, apolipoprotein AII, apolipoprotein B-100, lipoprotein apoCII, pre-apolipoprotein CIII, apolipoprotein apo C-IV, macrophage scavenger receptor type I, human antigen CD36 gene, serum amyloid P component, carboxyl ester lipase gene, paraoxonase 1, paraoxonase 2, paraoxonase 3, LDL-receptor related protein, hepatic triglyceride lipase, 3-hydroxy-3-methylglutaryl coenzyme A synthase, very low density lipoprotein receptor, microsomal triglyceride transfer protein, perilipin, endothelin-1, endothelin receptor A, interleukin 6, interleukin 1, complement protein C8 alpha, complement component C9, prostaglandin D2 synthase, annexin II/lipocortinII, annexin I/lipocortin, prostaglandin-endoperoxide synthase 2, insulin-like growth factor binding protein-1, secreted protein, acidic and rich in cysteine, human NF-kappa-B transcription factor, angiotensinogen, nitric oxide synthase 3, and nitric oxide synthase 2.

The invention also provides a substantially purified polynucleotide comprising a gene that is coexpressed with one or more known atherosclerosis-associated genes in a plurality of samples. In one aspect, the polynucleotide comprises a polynucleotide sequence selected from a polynucleotide encoding a peptide selected from SEQ ID NOs:1-34; a polynucleotide sequence complementary to the polynucleotide sequence of SEQ ID NOs:1-34; and a probe comprising at least 18 sequential nucleotides of the polynucleotide sequence of SEQ ID NOs:1-34 or their complements. The invention further provides a pharmaceutical composition comprising a polynucleotide and a pharmaceutical;

carrier.

The invention additionally provides methods for using a polynucleotide. One method uses the polynucleotide to screen a library of molecules or compounds to identify at least one ligand which specifically binds the polynucleotide and comprises combining the polynucleotide with a library of
5 molecules or compounds under conditions to allow specific binding and detecting specific binding, thereby identifying a ligand which specifically binds the polynucleotide. In this first method, the library is selected from DNA molecules, RNA molecules, PNAs, mimetics, and proteins; and the ligand identified using the method may be used to modulate the activity of the polynucleotide. A second
10 comprises combining the polynucleotide with a sample under conditions to allow specific binding, detecting specific binding between the polynucleotide and a ligand, recovering the bound polynucleotide, and separating the polynucleotide from the ligand, thereby obtaining purified ligand. A third method uses the polynucleotide to diagnose a disease or condition associated with the altered expression of a gene that is coexpressed with one or more known atherosclerosis-associated genes in a
15 plurality of biological samples and comprises hybridizing a polynucleotide to a sample under conditions to form one or more hybridization complexes, detecting the hybridization complexes, and comparing the levels of the hybridization complexes with the level of hybridization complexes in a non-diseased sample, wherein the altered level of hybridization complexes compared with the level of hybridization complexes of a non-diseased sample indicates the presence of the disease or condition.
20 A fourth method uses the polynucleotide to produce a polypeptide and comprises culturing a host cell containing an expression vector containing the polynucleotide under conditions for expression of the polypeptide and recovering the polypeptide from cell culture.

The invention provides a substantially purified polypeptide comprising the product of a gene that is coexpressed with one or more known atherosclerosis-associated genes in a plurality of samples.
25 The invention also provides a polypeptide comprising a polypeptide sequence selected from the polypeptides encoded by SEQ ID NOs:1-34 and an oligopeptide sequence comprising at least 6 sequential amino acids of the polypeptide sequence of encoded by SEQ ID NOs:1-34. The further provides a polypeptide comprising the amino acid sequence of SEQ ID NO:35. The invention still further provides a pharmaceutical composition comprising a polypeptide and a pharmaceutical carrier.

30 The invention additionally provides methods for using a polypeptide. One method uses the polypeptide to screen a library of molecules or compounds to identify at least one ligand which specifically binds the polypeptide and comprises combining the polypeptide with the library of molecules or compounds under conditions to allow specific binding and detecting specific binding between the polypeptide and ligand, thereby identifying a ligand which specifically binds the
35 polypeptide. In this method, the library is selected from DNA molecules, RNA molecules, PNAs,

mimetics, polypeptides, agonists, antagonists, and antibodies; and the ligand identified using the method is used to modulate the activity of the polypeptide. A second method uses the polypeptide to purify a ligand from a sample and comprises combining the polypeptide with a sample under conditions to allow specific binding, detecting specific binding between the polypeptide and a ligand, recovering the bound polypeptide, and separating the polypeptide from the ligand, thereby obtaining purified ligand. A third method uses the polypeptide to treat or to prevent a disease associated with the altered expression of a gene that is coexpressed with one or more known atherosclerosis-associated genes in a subject in need and comprises administering to the subject in need the pharmaceutical composition containing the polypeptide in an amount effective for treating the disease.

The invention provides an antibody or Fab comprising an antigen binding site, wherein the antigen binding site specifically binds to the polypeptide. The invention also provides a method for treating a disease associated with the altered expression of a gene that is coexpressed with one or more known atherosclerosis-associated genes in a subject in need, the method comprising the step of administering to the subject in need the antibody or the Fab in an amount effective for treating the disease. The invention further provides an immunoconjugate comprising the antigen binding site of the antibody or Fab joined to a therapeutic agent. The invention additionally provides a method for treating a disease associated with the altered expression of a gene that is coexpressed with one or more known atherosclerosis-associated genes in a subject in need, the method comprising the step of administering to the subject in need the immunoconjugate in an amount effective for treating the disease.

BRIEF DESCRIPTION OF THE SEQUENCE LISTING

The Sequence Listing provides exemplary atherosclerosis-associated gene sequences including polynucleotide sequences SEQ ID NOs:1-34 and the polypeptide sequence, SEQ ID NO:35. Each sequence is identified by a sequence identification number (SEQ ID NO).

DESCRIPTION OF THE INVENTION

It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the" include the plural reference unless the context clearly dictates otherwise. For example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Definitions

"Atherosclerosis-associated gene" refers to a gene or polynucleotide that exhibits a statistically significant coexpression pattern with known atherosclerosis-associated genes which are useful in the diagnosis, treatment, prognosis, or prevention of atherosclerosis.

"Known atherosclerosis-associated gene" refers to a sequence which has been previously identified as useful in the diagnosis, treatment, prognosis, or prevention of atherosclerosis and includes

polynucleotides encoding human 22kDa smooth muscle protein, calponin, desmin, smooth muscle myosin heavy chain, alpha tropomyosin, human tissue inhibitor of metalloproteinase 3, human tissue inhibitor of metalloproteinase-2, human tissue inhibitor of metalloproteinase-4, pro alpha 1(I) collagen, collagen alpha-2 type I, collagen alpha-6 type I, procollagen alpha 2(V), collagen VI alpha-2, type VI collagen alpha3, pro-alpha-1 type 3 collagen, pro-alpha-1 (V) collagen, collagenase type IV/ matrix metalloproteinase 9/gelatinase B, matrix Gla protein, cathepsin K, fibrinogen beta chain gene, fibrinogen gamma chain gene, pre-pro-von Willebrand factor, coagulation factor II/prothrombin, coagulation factor XII, coagulation factor VII, platelet endothelial cell adhesion molecule, lipoprotein-associated coagulation inhibitor, antithrombin III variant, plasminogen activator inhibitor-1, lipoprotein lipase, alpha-2-macroglobulin, apolipoprotein AI, apolipoprotein AII, apolipoprotein B-100, lipoprotein apoCII, pre-apolipoprotein CIII, apolipoprotein apo C-IV, macrophage scavenger receptor type I, human antigen CD36 gene, serum amyloid P component, carboxyl ester lipase gene, paraoxonase 1, paraoxonase 2, paraoxonase 3, LDL-receptor related protein, hepatic triglyceride lipase, 3-hydroxy-3-methylglutaryl coenzyme A synthase, very low density lipoprotein receptor, microsomal triglyceride transfer protein, perilipin, endothelin-1, endothelin receptor A, interleukin 6, interleukin 1, complement protein C8 alpha, complement component C9, prostaglandin D2 synthase, annexin II/lipocortinII, annexin I/lipocortin, prostaglandin-endoperoxide synthase 2, insulin-like growth factor binding protein-1, secreted protein, acidic and rich in cysteine, human NF-kappa-B transcription factor, angiotensinogen, nitric oxide synthase 3, and nitric oxide synthase 2. Typically, this means that the known gene is expressed at higher levels (i.e., has more abundant transcripts) in atherosclerotic lesions than in normal or non-diseased arterial intima or any other tissue.

"Ligand" refers to any molecule, agent, or compound which will bind specifically to a complementary site on a polynucleotide or polypeptide. Such ligands stabilize or modulate the activity of polynucleotides or polypeptides of the invention. For example, ligands are libraries of inorganic and organic molecules or compounds such as nucleic acids, proteins, peptides, carbohydrates, fats, and lipids.

"NSEQ" refers generally to a polynucleotide sequence of the present invention, including SEQ ID NO:1-34. "PSEQ" refers generally to a polypeptide sequence of the present invention, including SEQ ID NO:35.

A "fragment" refers to a nucleic acid sequence that is preferably at least 20 nucleotides in length, more preferably 40 nucleotides, and most preferably 60 nucleotides in length, and encompasses, for example, fragments consisting of 1-50, 51-400, 401-4000, 4001-12,000 nucleotides, and the like, of SEQ ID NO:1-34.

"Gene" refers to the partial or complete coding sequence of a gene including 5' or 3' untranslated regions. The gene may be in a sense or antisense (complementary) orientation.

"Polynucleotide" refers to a nucleic acid, nucleic acid sequence, oligonucleotide, nucleotide, or any fragment thereof. It may be DNA or RNA of genomic or synthetic origin, double-stranded or single-stranded, and combined with carbohydrate, lipids, protein or other materials to perform a particular activity or form a useful composition. "Oligonucleotide" is substantially equivalent to the
5 terms amplicon, primer, oligomer, element, and probe.

"Polypeptide" refers to an amino acid, amino acid sequence, oligopeptide, peptide, or protein or portions thereof whether naturally occurring or synthetic.

A "portion" refers to peptide sequence which is preferably at least 5 to about 15 amino acids in length, most preferably at least 10 amino acids long, and which retains some biological or
10 immunological activity of, for example, a portion of SEQ ID NO:35.

"Sample" is used in its broadest sense. A sample containing nucleic acids may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; genomic DNA, RNA, or cDNA in solution or bound to a substrate; a cell; a tissue; a tissue print; and the like.

"Substantially purified" refers to a nucleic acid or an amino acid sequence that is removed from
15 its natural environment and that is isolated or separated, and is at least about 60% free, preferably about 75% free, and most preferably about 90% free, from other components with which it is naturally present.

"Substrate" refers to any rigid or semi-rigid support to which polynucleotides or polypeptides are bound and includes membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic
20 beads, gels, capillaries or other tubing, plates, polymers, and microparticles with a variety of surface forms including wells, trenches, pins, channels and pores.

A "variant" refers to a polynucleotide or polypeptide whose sequence diverges from SEQ ID NO:1-35. Polynucleotide sequence divergence may result from mutational changes such as deletions, additions, and substitutions of one or more nucleotides; it may also be introduced to accommodate
25 differences in codon usage. Each of these types of changes may occur alone, or in combination, one or more times in a given sequence.

THE INVENTION

The present invention encompasses a method for identifying biomolecules that are associated with a specific disease, regulatory pathway, subcellular compartment, cell type, tissue type, or species.
30 In particular, the method identifies polynucleotides useful in diagnosis, prognosis, treatment, prevention, and evaluation of therapies for diseases associated with atherosclerosis including, but not limited to, stroke, myocardial infarction, hypertension, transient cerebral ischemia, mesenteric ischemia, coronary artery disease, angina pectoris, peripheral vascular disease, intermittent claudication, renal artery stenosis, and hypertension.

35 The method entails first identifying polynucleotides that are expressed in a plurality of cDNA

libraries. The identified polynucleotides include genes of known or unknown function which are expressed in a specific disease process, subcellular compartment, cell type, tissue type, or species. The expression patterns of the genes with known function are compared with those of genes with unknown function to determine whether a specified coexpression probability threshold is met. Through this comparison, a subset of the polynucleotides having a high coexpression probability with the known genes can be identified. The high coexpression probability correlates with a particular coexpression probability threshold which is preferably less than 0.001 and more preferably less than 0.00001.

The polynucleotides originate from cDNA libraries derived from a variety of sources including, but not limited to, eukaryotes such as human, mouse, rat, dog, monkey, plant, and yeast; prokaryotes such as bacteria; and viruses. These polynucleotides can also be selected from a variety of sequence types including, but not limited to, expressed sequence tags (ESTs), assembled polynucleotide sequences, full length gene coding regions, promoters, introns, enhancers, 5' untranslated regions, and 3' untranslated regions. To have statistically significant analytical results, the polynucleotides need to be expressed in at least three cDNA libraries.

The cDNA libraries used in the coexpression analysis of the present invention can be obtained from adrenal gland, biliary tract, bladder, blood cells, blood vessels, bone marrow, brain, bronchus, cartilage, chromaffin system, colon, connective tissue, cultured cells, embryonic stem cells, endocrine glands, epithelium, esophagus, fetus, ganglia, heart, hypothalamus, immune system, intestine, islets of Langerhans, kidney, larynx, liver, lung, lymph, muscles, neurons, ovary, pancreas, penis, peripheral nervous system, phagocytes, pituitary, placenta, pleurus, prostate, salivary glands, seminal vesicles, skeleton, spleen, stomach, testis, thymus, tongue, ureter, uterus, and the like. The number of cDNA libraries selected can range from as few as 3 to greater than 10,000. Preferably, the number of the cDNA libraries is greater than 500.

In a preferred embodiment, genes are assembled from related sequences, such as assembled sequence fragments derived from a single transcript. Assembly of the sequences can be performed using sequences of various types including, but not limited to, ESTs, extensions, or shotgun sequences. In a most preferred embodiment, the polynucleotide sequences are derived from human sequences that have been assembled using the algorithm disclosed in "Database and System for Storing, Comparing and Displaying Related Biomolecular Sequence Information", Lincoln *et al.* Serial No:60/079,469, filed March 26, 1998, incorporated herein by reference.

Experimentally, differential expression of the polynucleotides can be evaluated by methods including, but not limited to, differential display by spatial immobilization or by gel electrophoresis, genome mismatch scanning, representational difference analysis, and transcript imaging. Additionally, differential expression can be assessed by microarray technology. These methods may be used alone or in combination.

Known atherosclerosis-associated genes are selected based on the use of these genes as diagnostic or prognostic markers or as therapeutic targets.

The procedure for identifying novel genes that exhibit a statistically significant coexpression pattern with known atherosclerosis-associated genes is as follows. First, the presence or absence of a
 5 gene in a cDNA library is defined: a gene is present in a cDNA library when at least one cDNA fragment corresponding to that gene is detected in a cDNA sample taken from the library, and a gene is absent from a library when no corresponding cDNA fragment is detected in the sample.

Second, the significance of gene coexpression is evaluated using a probability method to measure a due-to-chance probability of the coexpression. The probability method can be the Fisher
 10 exact test, the chi-squared test, or the kappa test. These tests and examples of their applications are well known in the art and can be found in standard statistics texts (Agresti (1990) Categorical Data Analysis, John Wiley & Sons, New York NY; Rice (1988) Mathematical Statistics and Data Analysis, Duxbury Press, Pacific Grove CA). A Bonferroni correction (Rice, supra, p. 384) can also be applied in combination with one of the probability methods for correcting statistical results of one gene versus
 15 multiple other genes. In a preferred embodiment, the due-to-chance probability is measured by a Fisher exact test, and the threshold of the due-to-chance probability is set preferably to less than 0.001, more preferably to less than 0.00001. To determine whether two genes, A and B, have similar coexpression patterns, occurrence data vectors can be generated as illustrated in Table 1. The presence of a gene occurring at least once in a library is indicated by a one, and its absence from the library, by a
 20 zero.

Table 1. Occurrence data for genes A and B

	Library 1	Library 2	Library 3	...	Library N
gene A	1	1	0	...	0
gene B	1	0	1	...	0

For a given pair of genes, the occurrence data in Table 1 can be summarized in a 2 x 2 contingency
 table.

Table 2. Contingency table for co-occurrences of genes A and B

	Gene A present	Gene A absent	Total
Gene B present	8	2	10
Gene B absent	2	18	20
Total	10	20	30

Table 2 presents co-occurrence data for gene A and gene B in a total of 30 libraries. Both gene
 35 A and gene B occur 10 times in the libraries. Table 2 summarizes and presents: 1) the number of times

gene A and B are both present in a library; 2) the number of times gene A and B are both absent in a library; 3) the number of times gene A is present, and gene B is absent; and 4) the number of times gene B is present, and gene A is absent. The upper left entry is the number of times the two genes co-occur in a library, and the middle right entry is the number of times neither gene occurs in a library.

5 The off diagonal entries are the number of times one gene occurs, and the other does not. Both A and B are present eight times and absent 18 times. Gene A is present, and gene B is absent, two times; and gene B is present, and gene A is absent, two times. The probability ("p-value") that the above association occurs due to chance as calculated using a Fisher exact test is 0.0003. Associations are generally considered significant if a p-value is less than 0.01 (Agresti, *supra*; Rice, *supra*).

10 This method of estimating the probability for coexpression of two genes makes several assumptions. The method assumes that the libraries are independent and are identically sampled. However, in practical situations, the selected cDNA libraries are not entirely independent, because more than one library may be obtained from a single subject or tissue. Nor are they entirely identically sampled, because different numbers of cDNAs may be sequenced from each library. The number of
15 cDNAs sequenced typically ranges from 5,000 to 10,000 cDNAs per library. In addition, because a Fisher exact coexpression probability is calculated for each gene versus 45,233 other assembled genes, a Bonferroni correction for multiple statistical tests is used.

The present invention identifies 34 novel atherosclerosis-associated polynucleotides that exhibit strong association with genes known to be specific to atherosclerosis. The results presented in Table 4
20 show that the expression of the 34 novel atherosclerosis-associated polynucleotides has direct or indirect association with the expression of known atherosclerosis-associated genes. Therefore, the novel atherosclerosis-associated polynucleotides can potentially be used in diagnosis, treatment, prognosis, or prevention of diseases associated with atherosclerosis or in the evaluation of therapies for atherosclerosis. Further, the gene products of the 34 novel atherosclerosis-associated polynucleotides
25 are either potential therapeutics or targets of therapeutics against atherosclerosis.

Therefore, in one embodiment, the present invention encompasses a polynucleotide sequence comprising the sequence of SEQ ID NO:1-34. These 34 polynucleotides are shown by the method of the present invention to have strong coexpression association with known atherosclerosis-associated genes and with each other. The invention also encompasses a variant of the polynucleotide sequence,
30 its complement, or 18 consecutive nucleotides of a sequence provided in the above described sequences. Variant polynucleotide sequences typically have at least about 75%, more preferably at least about 85%, and most preferably at least about 95% polynucleotide sequence identity to NSEQ.

NSEQ or the encoded PSEQ may be used to search against the GenBank primate (pri), rodent (rod), mammalian (mam), vertebrate (vrtp), and eukaryote (eukp) databases, SwissProt, BLOCKS
35 (Bairoch *et al.* (1997) *Nucleic Acids Res* 25:217-221), PFAM, and other databases that contain

previously identified and annotated motifs, sequences, and gene functions. Methods that search for primary sequence patterns with secondary structure gap penalties (Smith *et al.* (1992) *Prot Eng* 5:35-51) as well as algorithms such as Basic Local Alignment Search Tool (BLAST; Altschul (1993) *J Mol Evol* 36:290-300; Altschul *et al.* (1990) *J Mol Biol* 215:403-410), BLOCKS (Henikoff and Henikoff (1991) *Nucleic Acids Res* 19:6565-6572), Hidden Markov Models (HMM; Eddy (1996) *Cur Opin Str Biol* 6:361-365; Sonnhammer *et al.* (1997) *Proteins* 28:405-420), and the like, can be used to manipulate and analyze nucleotide and amino acid sequences. These databases, algorithms and other methods are well known in the art and are described in Ausubel *et al.* (1997; Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7) and in Meyers (1995; Molecular Biology and Biotechnology, Wiley VCH, New York NY, p 856-853).

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to SEQ ID NO:1-34, and fragments thereof under stringent conditions. Stringent conditions can be defined by salt concentration, temperature, and other chemicals and conditions well known in the art. Conditions can be selected, for example, by varying the concentrations of salt in the prehybridization, hybridization, and wash solutions or by varying the hybridization and wash temperatures. With some substrates, the temperature can be decreased by adding formamide to the prehybridization and hybridization solutions.

Hybridization can be performed at low stringency, with buffers such as 5xSSC with 1% sodium dodecyl sulfate (SDS) at 60°C, which permits complex formation between two nucleic acid sequences that contain some mismatches. Subsequent washes are performed at higher stringency with buffers such as 0.2xSSC with 0.1% SDS at either 45°C (medium stringency) or 68°C (high stringency), to maintain hybridization of only those complexes that contain completely complementary sequences. Background signals can be reduced by the use of detergents such as SDS, Sarcosyl, or TRITON X-100 (Sigma-Aldrich, St. Louis MO), and/or a blocking agent, such as salmon sperm DNA. Hybridization methods are described in detail in Ausubel (*supra*, units 2.8-2.11, 3.18-3.19 and 4-6-4.9) and Sambrook *et al.* (1989; Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY)

NSEQ can be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences such as promoters and other regulatory elements. (See, e.g., Dieffenbach and Dveksler (1995) PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview NY). Additionally, one may use an XL-PCR kit (PE Biosystems, Foster City CA), nested primers, and commercially available cDNA libraries (Life Technologies, Rockville MD) or genomic libraries (Clontech, Palo Alto CA) to extend the sequence. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another program, to be about 18 to 30 nucleotides in length, to have a GC content of about 50%, and to form a hybridization complex at temperatures of

about 68°C to 72°C.

In another aspect of the invention, NSEQ can be cloned in recombinant DNA molecules that direct the expression of PSEQ, or structural or functional portions thereof, in host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express the polypeptide encoded by NSEQ. The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter the nucleotide sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In order to express a biologically active polypeptide, NSEQ, or derivatives thereof, may be inserted into an expression vector, i.e., a vector which contains the elements for transcriptional and translational control of the inserted coding sequence in a particular host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions. Methods which are well known to those skilled in the art may be used to construct such expression vectors. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, supra; and Ausubel, supra).

A variety of expression vector/host cell systems may be utilized to express NSEQ. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with baculovirus vectors; plant cell systems transformed with viral or bacterial expression vectors; or animal cell systems. For long term production of recombinant proteins in mammalian systems, stable expression in cell lines is preferred. For example, NSEQ can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable or visible marker gene on the same or on a separate vector. The invention is not to be limited by the vector or host cell employed.

In general, host cells that contain NSEQ and that express PSEQ may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or amino acid sequences. Immunological methods for detecting and measuring the expression of PSEQ using either specific polyclonal or monoclonal antibodies are known

in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS).

Host cells transformed with NSEQ may be cultured under conditions for the expression and recovery of the polypeptide from cell culture. The polypeptide produced by a transgenic cell may be
5 secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing NSEQ may be designed to contain signal sequences which direct secretion of the polypeptide through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the
10 inserted sequences or to process the expressed polypeptide in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the polypeptide may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational
15 activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the ATCC (Manassas VA) and may be chosen to ensure the correct modification and processing of the expressed polypeptide.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences are ligated to a heterologous sequence resulting in translation of a fusion polypeptide containing heterologous polypeptide moieties in any of the aforementioned host systems. Such
20 heterologous polypeptide moieties facilitate purification of fusion polypeptides using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase, maltose binding protein, thioredoxin, calmodulin binding peptide, 6-His, FLAG, *c-myc*, hemagglutinin, and monoclonal antibody epitopes.

In another embodiment, the nucleic acid sequences are synthesized, in whole or in part, using
25 chemical or enzymatic methods well known in the art (Caruthers *et al.* (1980) *Nucleic Acids Symp Ser* (7) 215-233; Ausubel, *supra*). For example, peptide synthesis can be performed using various solid-phase techniques (Roberge *et al.* (1995) *Science* 269:202-204), and machines such as the ABI 431A Peptide synthesizer (PE Biosystems) can be used to automate synthesis. If desired, the amino acid sequence may be altered during synthesis and/or combined with sequences from other proteins to
30 produce a variant protein.

In another embodiment, the invention entails a substantially purified polypeptide comprising the amino acid sequence of SEQ ID NO:35 and fragments thereof.

SCREENING, DIAGNOSTICS AND THERAPEUTICS

The polynucleotide sequences can be used in diagnosis, prognosis, treatment, prevention, and
35 selection and evaluation of therapies for atherosclerosis including, but not limited to, stroke, myocardial

infarction, hypertension, transient cerebral ischemia, mesenteric ischemia, coronary artery disease, angina pectoris, peripheral vascular disease, intermittent claudication, renal artery stenosis, and hypertension.

The polynucleotide sequences may be used to screen a library of molecules for specific binding
5 affinity. The assay can be used to screen a library of DNA molecules, RNA molecules, PNAs, peptides, ribozymes, antibodies, agonists, antagonists, immunoglobulins, inhibitors, proteins including transcription factors, enhancers, repressors, and drugs and the like which regulate the activity of the polynucleotide sequence in the biological system. The assay involves providing a library of molecules, combining the polynucleotide sequence or a fragment thereof with the library of molecules under
10 conditions suitable to allow specific binding, and detecting specific binding to identify at least one molecule which specifically binds the polynucleotide sequence.

Similarly the polypeptide or a portion thereof may be used to screen libraries of molecules in any of a variety of screening assays. The portion of the polypeptide employed in such screening may be free in solution, affixed to an abiotic or biotic substrate (e.g. borne on a cell surface), or located
15 intracellularly. Specific binding between the polypeptide and molecule may be measured. The assay can be used to screen a library of DNA molecules, RNA molecules, PNAs, peptides, mimetics, ribozymes, antibodies, agonists, antagonists, immunoglobulins, inhibitors, peptides, polypeptides, drugs and the like, which specifically bind the polypeptide. One method for high throughput screening using very small assay volumes and very small amounts of test compound is described in Burbaum et
20 al. USPN 5,876,946, incorporated herein by reference, which screens large numbers of molecules for enzyme inhibition or receptor binding.

In one preferred embodiment, the polynucleotide sequences are used for diagnostic purposes to determine the absence, presence, and excess expression of the polypeptide. The polynucleotides may be at least 18 nucleotides long and consist of complementary RNA and DNA molecules, branched
25 nucleic acids, and/or peptide nucleic acids (PNAs). In one alternative, the polynucleotides are used to detect and quantify gene expression in samples in which expression of NSEQ is correlated with disease. In another alternative, NSEQ can be used to detect genetic polymorphisms associated with a disease. These polymorphisms may be detected in the transcript cDNA.

The specificity of the probe is determined by whether it is made from a unique region, a
30 regulatory region, or from a conserved motif. Both probe specificity and the stringency of diagnostic hybridization or amplification (maximal, high, intermediate, or low) will determine whether the probe identifies only naturally occurring, exactly complementary sequences, allelic variants, or related sequences. Probes designed to detect related sequences should preferably have at least 75% sequence identity to any of the polynucleotides encoding PSEQ.

35 Methods for producing hybridization probes include the cloning of nucleic acid sequences into

vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by adding RNA polymerases and labeled nucleotides. Hybridization probes may incorporate nucleotides labeled by a variety of reporter groups including, but not limited to, radionuclides such as ^{32}P or ^{35}S , enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, fluorescent labels, and the like. The labeled polynucleotide sequences may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; and in microarrays utilizing samples from subjects to detect altered PSEQ expression.

NSEQ can be labeled by standard methods and added to a sample from a subject under conditions for the formation and detection of hybridization complexes. After incubation the sample is washed, and the signal associated with hybrid complex formation is quantitated and compared with a standard value. Standard values are derived from any control sample, typically one that is free of the suspect disease. If the amount of signal in the subject sample is altered in comparison to the standard value, then the presence of altered levels of expression in the sample indicates the presence of the disease. Qualitative and quantitative methods for comparing the hybridization complexes formed in subject samples with previously established standards are well known in the art.

Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual subject. Once the presence of disease is established and a treatment protocol is initiated, hybridization or amplification assays can be repeated on a regular basis to determine if the level of expression in the subject begins to approximate that which is observed in a healthy subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to many years.

The polynucleotides may be used for the diagnosis of a variety of diseases associated with atherosclerosis. These include, but are not limited to, stroke, myocardial infarction, hypertension, transient cerebral ischemia, mesenteric ischemia, coronary artery disease, angina pectoris, peripheral vascular disease, intermittent claudication, renal artery stenosis, and hypertension.

The polynucleotides may also be used as targets in a microarray. The microarray can be used to monitor the expression patterns of large numbers of genes simultaneously and to identify splice variants, mutations, and polymorphisms. Information derived from analyses of the expression patterns may be used to determine gene function, to understand the genetic basis of a disease, to diagnose a disease, and to develop and monitor the activities of therapeutic agents used to treat a disease. Microarrays may also be used to detect genetic diversity, single nucleotide polymorphisms which may characterize a particular population, at the genome level.

In yet another alternative, polynucleotides may be used to generate hybridization probes useful

in mapping the naturally occurring genomic sequence. Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data as described in Heinz-Ulrich et al. (In: Meyers, supra, pp. 965-968).

In another embodiment, antibodies or Fabs comprising an antigen binding site that specifically
5 binds PSEQ may be used for the diagnosis of diseases characterized by the over-or-under expression of PSEQ. A variety of protocols for measuring PSEQ, including ELISAs, RIAs, and FACS, are well known in the art and provide a basis for diagnosing altered or abnormal levels of expression. Standard values for PSEQ expression are established by combining samples taken from healthy subjects, preferably human, with antibody to PSEQ under conditions for complex formation. The amount of
10 complex formation may be quantitated by various methods, preferably by photometric means. Quantities of PSEQ expressed in disease samples are compared with standard values. Deviation between standard and subject values establishes the parameters for diagnosing or monitoring disease. Alternatively, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PSEQ specifically compete with a test compound for binding the polypeptide. Antibodies
15 can be used to detect the presence of any peptide which shares one or more antigenic determinants with PSEQ. In one aspect, the anti-PSEQ antibodies of the present invention can be used for treatment or monitoring therapeutic treatment for atherosclerosis.

In another aspect, the NSEQ, or its complement, may be used therapeutically for the purpose of expressing mRNA and polypeptide, or conversely to block transcription or translation of the mRNA.
20 Expression vectors may be constructed using elements from retroviruses, adenoviruses, herpes or vaccinia viruses, or bacterial plasmids, and the like. These vectors may be used for delivery of nucleotide sequences to a particular target organ, tissue, or cell population. Methods well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences or their complements. (See, e.g., Maulik et al. (1997) Molecular Biotechnology, Therapeutic Applications and
25 Strategies, Wiley-Liss, New York NY.) Alternatively, NSEQ, or its complement, may be used for somatic cell or stem cell gene therapy. Vectors may be introduced in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors are introduced into stem cells taken from the subject, and the resulting transgenic cells are clonally propagated for autologous transplant back into that same subject. Delivery of NSEQ by transfection, liposome injections, or polycationic amino polymers may be achieved using methods
30 which are well known in the art and described in Goldman et al. (1997; *Nature Biotechnol* 15:462-466). Additionally, endogenous NSEQ expression may be inactivated using homologous recombination methods which insert an inactive gene sequence into the coding region or other targeted region of NSEQ. (See, e.g. Thomas et al. (1987) *Cell* 51:503-512.)

Vectors containing NSEQ can be transformed into a cell or tissue to express a missing
35 polypeptide or to replace a nonfunctional polypeptide. Similarly a vector constructed to express the

complement of NSEQ can be transformed into a cell to downregulate the overexpression of PSEQ. Complementary or antisense sequences may consist of an oligonucleotide derived from the transcription initiation site; nucleotides between about positions -10 and +10 from the ATG are preferred. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful
5 because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee *et al.* In: Huber and Carr (1994) Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco NY, pp. 163-177.)

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the cleavage of mRNA
10 and decrease the levels of particular mRNAs, such as those comprising the polynucleotide sequences of the invention. (See, e.g., Rossi (1994) *Current Biology* 4: 469-471.) Ribozymes may cleave mRNA at specific cleavage sites. Alternatively, ribozymes may cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The construction and production of ribozymes is well known in the art and is described in Meyers (*supra*).

15 RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiester linkages within the backbone of the molecule. Alternatively, nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine,
20 guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases, may be included.

Further, an antagonist, or an antibody that binds specifically to PSEQ may be administered to a subject to treat or prevent atherosclerosis. The antagonist, antibody, or fragment may be used directly to inhibit the activity of the polypeptide or indirectly to deliver a therapeutic agent to cells or tissues
25 which express the PSEQ. An immunoconjugate comprising a PSEQ binding site of the antibody or the antagonist and a therapeutic agent may be administered to a subject in need to treat or prevent disease.

The therapeutic agent may be a cytotoxic agent selected from a group including, but not limited to, abrin, ricin, doxorubicin, daunorubicin, taxol, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin D, diphtheria toxin,

30 Pseudomonas exotoxin A and 40, radioisotopes, and glucocorticoid.

Antibodies to PSEQ may be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies, such as those which inhibit dimer formation, are especially preferred for therapeutic use.

35 Monoclonal antibodies to PSEQ may be prepared using any technique which provides for the

production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma, the human B-cell hybridoma, and the EBV-hybridoma techniques. In addition, techniques developed for the production of chimeric antibodies can be used. (See, e.g., Pound (1998) Immunochemical Protocols, Methods Mol Biol Vol 80). Alternatively, techniques described for the
5 production of single chain antibodies may be employed. Fabs which contain specific binding sites for PSEQ may also be generated. Various immunoassays may be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art.

Yet further, an agonist of PSEQ may be administered to a subject to treat or prevent a disease
10 associated with decreased expression, longevity or activity of PSEQ.

An additional aspect of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic applications discussed above. Such pharmaceutical compositions may consist of PSEQ or antibodies, mimetics, agonists, antagonists, or inhibitors of the polypeptide. The compositions may be
15 administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a subject alone or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number
20 of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing
25 of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also
30 be used to determine the concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating and
35 contrasting the ED₅₀ (the dose therapeutically effective in 50% of the population) and LD₅₀ (the dose

lethal to 50% of the population) statistics. Any of the therapeutic compositions described above may be applied to any subject in need of such therapy, including, but not limited to, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

EXAMPLES

5 It is to be understood that this invention is not limited to the particular devices, machines, materials and methods described. Although particular embodiments are described, equivalent embodiments may be used to practice the invention. The described embodiments are provided to illustrate the invention and are not intended to limit the scope of the invention which is limited only by the appended claims.

10 I cDNA Library Construction

The cDNA library SMCCNOS01 was selected as an example to demonstrate the construction of cDNA libraries from which the polynucleotides associated with known atherosclerosis-associated genes were derived. The SMCCNOS01 subtracted coronary artery smooth muscle cell library was constructed using 7.56×10^6 clones from the SMCCNOT02 library and was subjected to two rounds of
15 subtraction hybridization for 48 hours with 6.12×10^6 clones from SMCCNOT01. The SMCCNOT02 library was constructed using RNA isolated from coronary artery smooth muscle cells removed from a 3-year-old Caucasian male. The cells were treated for 20 hours with $\text{TNF}\alpha$ and $\text{IL-1}\beta$ at 10ng/ml each. The SMCCNOT01 was constructed using RNA isolated from untreated coronary artery smooth muscle cells from the same donor. Subtractive hybridization conditions were based on the methodologies of
20 Swaroop *et al.* (1991; Nucleic Acids Res 19:1954) and Bonaldo *et al.* (1996; Genome Res 6:791).

For both cDNA libraries, SMCCNOT01 and SMCCNOT02, the frozen coronary artery smooth muscle cells (50-100 mg) were homogenized in GTC buffer (4.0M guanidine thiocyanate, 0.1M Tris-HCl pH 7.5, 1% 2-mercaptoethanol). Two volumes of binding buffer (0.4M LiCl, 0.1M Tris-HCl pH 7.5, 0.02M EDTA) were added, and the resulting mixture was vortexed at 13,000 rpm. The supernatant
25 was removed and combined with Oligo d(T)₂₅ bound streptavidin particles (MPG). After rotation at room temperature, the mRNA-Oligo d(T)₂₅ bound streptavidin particles were separated from the supernatant, washed twice with hybridization buffer I (0.15M NaCl, 0.01M Tris-HCl pH8.0, 1mM EDTA, 0.1% lauryl sarcosinate) using magnetic separation at each step to remove the supernatant from the particles. Bound mRNA was eluted from the particles with release solution and heated to 65°C.
30 The supernatant containing eluted mRNA was magnetically separated from the particles and used to construct the cDNA libraries.

The RNA was used for cDNA synthesis and construction of the cDNA library according to the recommended protocols in the SUPERScript plasmid system (Life Technologies). The cDNAs were fractionated on a SEPHAROSE CL4B column (Amersham Pharmacia Biotech (APB), Piscataway NJ),
35 and those cDNAs exceeding 400 bp were ligated into pINCY plasmid (Incyte Genomics, Palo Alto

CA). Recombinant plasmids were transformed into DH5 α competent cells or ELECTROMAX cells (Life Technologies).

II Isolation and Sequencing of cDNA Clones

Plasmid DNA was released from the cells and purified using the REAL PREP 96 plasmid kit (Qiagen, Valencia CA). The recommended protocol was employed except for the following changes:
1) the bacteria were cultured in 1 ml of sterile TERRIFIC BROTH media (BD Biosciences, Sparks MD) with carbenicillin at 25 mg/L and glycerol at 0.4%; 2) after inoculation, the cells were cultured for 19 hours and then lysed with 0.3 ml of lysis buffer; and 3) following isopropanol precipitation, the plasmid DNA pellet was resuspended in 0.1 ml distilled water, and samples were transferred to a 96-well block for storage at 4° C.

The cDNAs were prepared using a MICROLAB 2200 System (Hamilton, Reno NV) in combination with the DNA ENGINE thermal cycler (MJ Research, Watertown MA). cDNAs were sequenced by the method of Sanger and Coulson (1975; J Mol Biol 94:441f) using ABI PRISM 377 (PE Biosystems) or MEGABACE 1000 sequencing systems (APB).

Most of the sequences disclosed herein were sequenced using standard ABI protocols and kits (PE Biosystems) at solution volumes of 0.25x -1.0x concentrations. In the alternative, some of the sequences disclosed herein were sequenced using solutions and dyes from APB.

III Selection, Assembly, and Characterization of Sequences

The sequences used for co-expression analysis were assembled from EST sequences, 5' and 3' longread sequences, and full length coding sequences. Selected assembled sequences were expressed in at least three cDNA libraries.

The assembly process is described as follows. EST sequence chromatograms were processed and verified. Quality scores were obtained using PHRED (Ewing *et al.* (1998) Genome Res 8:175-185; Ewing and Green (1998) Genome Res 8:186-194), and edited sequences were loaded into a relational database management system (RDBMS). The sequences were clustered using BLAST with a product score of 50. All clusters of two or more sequences created a bin which represents one transcribed gene.

Assembly of the component sequences within each bin was performed using a modification of Phrap, a publicly available program for assembling DNA fragments (Green, P. University of Washington, Seattle WA). Bins that showed 82% identity from a local pair-wise alignment between any of the consensus sequences were merged.

Bins were annotated by screening the consensus sequence in each bin against public databases, such as GBpri and GenPept from NCBI. The annotation process involved a FASTn screen against the GBpri database in GenBank. Those hits with a percent identity of greater than or equal to 75% and an alignment length of greater than or equal to 100 base pairs were recorded as homolog hits. The residual unannotated sequences were screened by FASTx against GenPept. Those hits with an E value of less

than or equal to 10^{-8} were recorded as homolog hits.

Sequences were then reclustered using BLASTn and Cross-Match, a program for rapid amino acid and nucleic acid sequence comparison and database search (Green, *supra*), sequentially. Any BLAST alignment between a sequence and a consensus sequence with a score greater than 150 was realigned using cross-match. The sequence was added to the bin whose consensus sequence gave the highest Smith-Waterman score (Smith *et al.* (1992) *Incyte Genomics* 5:35-51) amongst local alignments with at least 82% identity. Non-matching sequences were moved into new bins, and assembly processes were repeated.

IV Coexpression Analyses of Atherosclerosis-Associated Genes

Sixty-six known atherosclerosis-associated genes were selected to identify novel genes that are closely associated with atherosclerosis. The known atherosclerosis-associated genes which were examined in this analysis and brief descriptions of their functions are listed in Table 3.

Table 3. Descriptions of Known Atherosclerosis-Associated Genes

	GENE	DESCRIPTION AND REFERENCES
15	Human 22kDa smooth muscle protein (SM22)	Smooth muscle cell-specific gene which is down-regulated during smooth muscle cell dedifferentiation as part of atherogenic process (Sobue <i>et al.</i> (1998) <i>Horm Res</i> 50:15-24; Sobue <i>et al.</i> (1999) <i>Mol Cell Biochem</i> 190:105-18)
	calponin (CNN1)	Calponin is smooth muscle-specific and may mediate smooth muscle contractility through it's binding of the amino-terminal end of the myosin regulatory light chain. Involved in phenotypic modulation of smooth muscle cells, a feature of atherosclerosis (Szymanski <i>et al.</i> (1999) <i>Biochemistry</i> 38:3778-84)
20	desmin (DES)	Contractile component of myofibrils in differentiated smooth muscle cells. Regarded as a marker for smooth muscle cells (Shi <i>et al.</i> (1997) <i>Circulation</i> 95:2684-93)
	smooth muscle myosin heavy chain (MYH11)	Contractile component of myofibrils in differentiated smooth muscle cells. Regarded as a marker for smooth muscle cells (Sobue <i>et al.</i> (1999) <i>Mol Cell Biochem</i> 190:105-18)
25	alpha tropomyosin (TPM1)	Contractile component of myofibrils in differentiated smooth muscle cells (Sobue <i>et al.</i> (1999) <i>Mol Cell Biochem</i> 190:105-18; Kashiwada <i>et al.</i> (1997) <i>J Biol Chem</i> 272:15396-404)
30	Human tissue inhibitor of metalloproteinase 3 (TIMP3)	TIMPs control the activity of matrix metalloproteinases and are important in local matrix remodeling of vasculature. Atheroma extracts shown to have 5x higher TIMP3 expression levels than non-atherosclerotic tissue. Abundant TIMP1,2, 3 expression noted in plaque macrophages and smooth muscle cells. PDGF and TGFbeta augment TIMP3 expression. TIMP3 possible important role in plaque stability (Fabunmi <i>et al.</i> (1998) <i>Circ Res</i> 83:270-8)

	Human tissue inhibitor of metalloproteinase-2 (TIMP-2)	TIMPs control the activity of matrix metalloproteinases and are important in local matrix remodeling of vasculature. Abundant TIMP1,2, 3 expression noted in plaque macrophages and smooth muscle cells. Expression of TIMP2 is greatly increased during neointima formation in organ cultures of human saphenous vein (Kranzhofer <i>et al.</i> (1999) <i>Arterioscler Thromb Vasc Biol</i> 19:255-65)
5	Human tissue inhibitor of metalloproteinase-4 (TIMP4)	TIMPs control the activity of matrix metalloproteinases and are important in local matrix remodeling of vasculature (Greene <i>et al.</i> (1996) <i>J Biol Chem</i> 271:30375-80)
10	pro alpha 1(I) collagen (COL1A1)	Member of family of fibrous structural proteins. Most abundant structural component of the extracellular matrix. Secreted as procollagen and converted to collagen by matrix metalloproteinases. Collagens are important in atherosclerosis for promoting platelet aggregation and for providing sites for platelet adhesion to the vessel wall (Wen <i>et al.</i> (1999) <i>Arterioscler Thromb Vasc Biol</i> 19:519-24)
	collagen alpha-2 type I (COL1A2)	see COL1A1 above
	COL6A1	see COL1A1 above
15	procollagen alpha 2(V) (COL5A2)	see COL1A1 above
	collagen VI alpha-2 (COL6A2)	see COL1A1 above
20	type VI collagen alpha3 (COL6A3)	see COL1A1 above
	pro-alpha-1 type 3 collagen (COL3A1)	see COL1A1 above
25	pro-alpha-1 (V) collagen (COL3A1)	see COL1A1 above
30	collagenase type IV/ matrix metalloproteinase 9/gelatinase B (MMP9)	Contributes to the degradation of vascular wall/smooth muscle cells associated with local matrix remodeling. Expression of metalloproteinases controlled by tissue inhibitors of metalloproteinases (TIMPs). Balance between MMP and TIMP expression becomes distorted during onset and progression of atherosclerosis. MMP9 localized to lesional macrophages, along with MMP-1, MMP-2, MMP-3. Rabbit aortic macrophage foam cells express immunoreactive MMP-9 (Moreau <i>et al.</i> (1999) <i>Circulation</i> 99:420-426; Zaltsman <i>et al.</i> (1997) <i>Atherosclerosis</i> 130:61-70)
	matrix Gla protein (MGP)	Role in active calcification of vascular smooth muscle cells, suggested by expression study on VSMC <i>in vitro</i> differentiation study. Calcifying phenotype associated with high MGP levels. MGP knockout mice develop to term, but die up to 2 months after birth due to extensive calcification of the arteries, causing blood vessel rupture (Luo <i>et al.</i> (1997) <i>Nature</i> 386:78-81; Mori <i>et al.</i> (1998) <i>FEBS Lett</i> 433:19-22)

	cathepsin K (CTSK)	Nonmetalloenzyme, potent elastase present in advanced atherosclerotic plaques. Contributes to the breakdown of components of vascular extracellular matrix, reducing tensile strength, increasing plaque vulnerability (Sukhova <i>et al.</i> (1998) <i>J Clin Invest</i> 102:576-83)
	fibrinogen beta chain gene (FGB)	Component of fibrin in the extracellular matrix. Fibrin deposition is an integral part of advanced atherosclerotic lesion development. Variation at the beta fibrinogen locus associated with peripheral atherosclerosis (Sueishi <i>et al.</i> (1998) <i>Semin Thromb Hemost</i> 24:255-260; Fowkes <i>et al.</i> (1992) <i>Lancet</i> 339:693-696)
5	fibrinogen beta chain gene (FGG)	Participant in adhesion and aggregation of platelets which occurs through binding of platelet receptors. FGG carries the main binding site for the platelet receptor binding. Mutations in FGG associated with clotting defects and thrombotic tendency. Fibrin deposition is an integral part of advanced atherosclerotic lesion development (Sueishi <i>et al.</i> (1998) <i>Semin Thromb Hemost</i> 24:255-60; Cote <i>et al.</i> (1998) <i>Blood</i> 92:2195-2212)
	pre-pro-von Willebrand factor (VWF)	Blood glycoprotein involved in normal hemostasis. Mediates adhesion of platelets to sites of vascular damage. Also acts as a cofactor in factor VIII activity in blood coagulation. Increased levels of VWF are found in atherosclerosis and in several of its major risk factors, including hypercholesterolemia, diabetes, obesity, hypertension. Levels serve as a predictor of adverse clinical outcome following vascular surgery, possibly as an indicator of thrombus formation (Sadler (1998) <i>Annu Rev Biochem</i> 67:395-424; Blann <i>et al.</i> (1994) <i>Eur J Vasc Surg</i> 8:10-15; Kessler <i>et al.</i> (1998) <i>Diabetes Metab</i> 24:327-36; Folsom <i>et al.</i> (1997) <i>Circulation</i> 96:1102-1108)
10	coagulation factor II/ prothrombin (F2)	Central role in blood hemostasis by regulating platelet aggregation and blood coagulation. Converts fibrinogen to fibrin in the final stage of clotting cascade. Promotes cellular chemotaxis and proliferation, extracellular matrix turnover and release of inflammatory cytokines (Goldsack <i>et al.</i> (1998) <i>Int J Biochem Cell Biol</i> 30:641-646)
	coagulation factor XII (F12)	Activation of blood coagulation is an important part of post-vascular injury with initiation of atherosclerotic lesion formation and contributes to thrombosis in advanced stage atherosclerosis (Sueishi <i>et al.</i> (1998) <i>Semin Thromb Hemost</i> 24:255-260)
15	coagulation factor VII (F7)	Central role in coagulation, influences plasma triglyceride levels, a risk factor in atherosclerosis. Epidemiological studies have linked F7 with cardiovascular risk/ atherothrombotic tendency (Ghaddar <i>et al.</i> (1998) <i>Circulation</i> 98:2815-2821; Koenig (1998) <i>Eur Heart J</i> 19:C39-43; Folsom <i>et al.</i> (1997) <i>Circulation</i> 96:1102-1108)
20	platelet endothelial cell adhesion molecule (PECAM-1)	Signalling molecule in the migration of cells as part of the pathophysiology of vascular occlusive diseases such as atherosclerosis. Analysis of endothelial/monocyte co-cultures indicates oxidative stress induces transendothelial migration of monocytes as a result of phosphorylation of PECAM-1 (Rattan <i>et al.</i> (1997) <i>Am J Physiol</i> 273:E453-61)

	lipoprotein-associated coagulation inhibitor (LACI)	Natural anticoagulant, inhibits factor VII/tissue factor complexes. Role in regulating coagulation in atherosclerotic plaques. Circulates in association with plasma lipoproteins VLDL, HDL and LDL. <u>In situ</u> expression studies indicate TFPI is expressed in adventitial layer of large arteries, and in atherosclerotic vessels is expressed by macrophages in focal areas throughout the plaque (Drew <u>et al.</u> (1997) Lab Invest 77:291-298; Sandset (1996) Haemostasis 26:154-165)
5	antithrombin III variant (AT3)	ATIII is the sole blood component through which heparin exerts its anti-coagulation effect. Deficiency in ATIII causes recurrent venous thrombosis and pulmonary embolism and can be inherited in autosomal dominant fashion (Hultin <u>et al.</u> (1988) Thromb Haemost 59:468-73; Lane <u>et al.</u> (1996) Blood Rev 10:59-74)
10	plasminogen activator inhibitor-1 (PAI-1)	Major physiological inhibitor of fibrinolysis. Plasma levels correlate with incidence of MI and venous thrombosis. Both adipocytes and endothelial cells produced PAI, possibly under the control of PPARG, as demonstrated using recombinant PPARG expression constructs in endothelial cell lines. Increased expression of PAI observed in coronary heart disease. 4G polymorphism in promotor causes increased PAI expression associated with MI in some studies (Eriksson <u>et al.</u> (1995) Proc Natl Acad Sci 92:1851-5; Marx <u>et al.</u> (1999) Arterioscler Thromb Vasc Biol 19:546-551)
	lipoprotein lipase (LPL)	Hydrolises triglyceride in chylomicrons and therefore regulates metabolism of circulating lipoproteins. Appears to have an atherogenic effect on the arterial wall due to its ability to alter the properties of LDL. Increased activity of LPL is found in atherosclerotic arteries when compared to normal. Expressed by macrophages in atherosclerotic lesions. Mutations in LPL responsible for familial hypercholesterolemia and premature atherosclerosis (Fisher <u>et al.</u> (1997) Atherosclerosis 135:145-159; Goldberg (1996) J Lipid Res 37:693-707; Gerdes <u>et al.</u> (1997) Circulation 96:733-740)
15	alpha-2-macroglobulin (A2M)	Foam cell formation - retains LDL cholesterol in the lipid core of atherosclerotic plaque (Llorente <u>et al.</u> (1998). Rev Esp Cardiol 51:633-641)
	apolipoprotein AI (APOA1)	Participates in reverse cholesterol transport from tissues to the liver. Promotes cholesterol efflux from tissues and acts as a cofactor for lecithin cholesterol acyltransferase (LCAT). Mutations in ApoA1 and of ApoA1/CIII/AIV gene cluster assoc with atherosclerosis. Transgenic mice expressing high plasma APOA1 levels are protected from fatty streak development with a high atherogenic diet (Gordon <u>et al.</u> (1989) Circulation 79:8-15; Rubin <u>et al.</u> (1991) Nature 353:265-7; Karathanasis <u>et al.</u> (1987) Proc Natl Acad Sci 84:7198-7202)
	apolipoprotein AII (APOA2)	Major component of HDL. Appears to have an opposite effect to that of APOAI, though exact function unknown. APOAII may have ability to convert HDL from an anti- to a pro-inflammatory particle, with paraoxonase having a role in this transformation process. Plasma APOAII levels significantly associated with plasma free fatty acid levels. Transgenic mice expressing varying levels of APOAII show increased atherosclerotic lesions than wt when fed an atherogenic diet. Possible interaction between diet/genotype and atherogenic potential (Escola-Gil <u>et al.</u> (1998) J Lipid Res 39:457-462; Warden <u>et al.</u> (1993) Proc Natl Acad Sci 90:10886-10890)

	apolipoprotein B-100 (APOB)	Main apolipoprotein of chylomicrons and low density lipoproteins. Mutations in APOB100 underly familial defective apolipoprotein B-100 in which patients suffer from premature atherosclerosis. Mutations result in defect in binding of LDL to LDL receptor, and accumulation of plasma LDL. High-expressing APOB transgenic mice exhibit elevated VLDL-LDL cholesterol and atherogenic lesions (Callow <i>et al.</i> (1995) J Clin Invest 96:1639-1646; Brasaemle <i>et al.</i> (1997) J Biol Chem 272:9378-9387)
	lipoprotein apoCII (APOC2)	Role in lipoprotein metabolism. Cofactor in the activity of lipoprotein lipase the enzyme that hydrolyzes triglycerides in plasma and transfers the fatty acids to tissues. Mutations in APOC2 responsible for hyperlipoproteinemia 1B, similar to lipoprotein lipase deficiency (Cox <i>et al.</i> (1978) N Engl J Med 299:1421-1424; Arimoto <i>et al.</i> (1998) J Lipid Res 39:143-151)
5	pre-apolipoprotein CIII (APOC3)	Inhibits lipoprotein lipase and hepatic lipase, decreases uptake of lymph chylomicrons by hepatic cells. APOA3 possibly delays breakdown of triglyceride rich particles. SstI RFLP in apoCIII is associated with plasma triglyceride and apoCIII levels and hyperlipidemic phenotypes (Henderson <i>et al.</i> (1987) Hum Genet 75:62-65)
	apolipoprotein apoC-IV (APOC4)	APOC4 is a lipid-binding protein that has the potential to alter lipid metabolism. Human APOC4 transgenic mice are hypertriglyceridaemic compared to normal controls (Allan <i>et al.</i> (1996) J Lipid Res 37:1510-1518)
10	macrophage scavenger receptor type I (MSR1)	Mediates binding, internalisation and processing of negatively-charged macromolecules. Implicated in the pathological deposition of cholesterol in arterial walls during atherogenesis (Han <i>et al.</i> (1998) Hum Mol Genet 7:1039-1046)
15	Human antigen CD36 gene (CD36)	Acts as a scavenger receptor for oxidised LDL. Transient regulation under control of M-CSF during monocyte-macrophage differentiation increases foam cell accumulation. Possible role in atherogenesis: increased M-CSF levels detected in atherosclerotic lesions in rabbits and humans. (Huh <i>et al.</i> (1996) Blood 87:2020-2028; Aitman <i>et al.</i> (1999) Nat Genet 21:76-83)
	serum amyloid P component (SAP)	Plasma glycoprotein expressed in atherosclerotic lesions. Interacts with lipoproteins in specific manner (Li <i>et al.</i> (1995) Arterioscler Thromb Vasc Biol 15:252-257; Li <i>et al.</i> (1998) Biochem Biophys Res Commun 244:249-252)
	carboxyl ester lipase gene (CEL)	CEL gene expression increases in presence of oxidised and native LDL <i>in vitro</i> . It is expressed in the vessel wall and in aortic extracts - may interact with cholesterol to modulate progression of atherosclerosis (Li <i>et al.</i> (1998) Biochem J 329:675-679)
20	paraoxonase 1 (PON1)	Serum esterase exclusively associated with high-density lipoproteins; it might confer protection against coronary artery disease by destroying pro-inflammatory oxidized lipids in oxidized low-density lipoproteins. PON1 gln192-to-arg polymorphism associated with CAD. Association between PON1 genetic variation and plasma LDL, HDL and non-HDL and apoB levels in genetically isolated Alberta Hutterite population. When fed on a high-fat, high-cholesterol diet, PON1-null mice were more susceptible to atherosclerosis than wild-type (Serrato <i>et al.</i> (1995) J Clin Invest 96:3005-3008; Boright <i>et al.</i> (1998) Atherosclerosis 139:131-136; Shih <i>et al.</i> (1998) Nature 394:284-287)

	paraoxonase 2 (PON2)	Serum esterase exclusively associated with high-density lipoproteins; it might confer protection against coronary artery disease by destroying pro-inflammatory oxidized lipids in oxidized low-density lipoproteins. Common polymorphism at codon 311 (cys-ser) in PON2 associated with CHD alone and synergistically with the 192 polymorphism in PON1 in Asian Indians. Association between genetic variation in PON2 and plasma cholesterol and apolipoprotein A1 in genetically isolated Alberta Hutterite population (Sanghera <i>et al.</i> (1998) <i>Am J Hum Genet</i> 62:36-44; Boright <i>et al.</i> (1998) <i>Atherosclerosis</i> 139:131-136)
	paraoxonase 3 (PON3)	Serum esterase exclusively associated with high-density lipoproteins; it might confer protection against coronary artery disease by destroying pro-inflammatory oxidized lipids in oxidized low-density lipoproteins. Other members PON2, 3 associated with CHD and cholesterol levels (Laplaud <i>et al.</i> (1998) <i>Clin Chem Lab Med</i> 36:431-441)
5	LDL-receptor related protein (LRP1)	Possible important role in atherosclerotic lesion development. Abundant expression of mRNA and protein found in vascular smooth muscle cells and macrophages of early and advanced atherosclerotic lesions. Receptor for uptake of ApoE-containing lipoprotein particles (Beisiegel <i>et al.</i> (1989) <i>Nature</i> 341:162-164; Hiltunen <i>et al.</i> (1998) <i>Atherosclerosis</i> 137:S81-88)
10	hepatic triglyceride lipase (HTGL)	Hepatic lipase is involved in cholesterol efflux. Downstream of cholesterol ester transfer protein in pathway: acts on triglyceride-rich HDL to promote formation of smaller HDL particles - effectors of cellular cholesterol efflux (Fan <i>et al.</i> (1998) <i>J Atheroscler Thromb</i> 5:41-45; Santamarina-Fojo <i>et al.</i> (1998) <i>Curr Opin Lipidol</i> 9:211-219)
15	3-hydroxy-3-methylglutaryl coenzyme A synthase (HMGCR)	Catalyses rate limiting step in cholesterol biosynthesis as well as being involved in other systems (eg. primordial germ cell migration). Expression of HMG CoA reductase is regulated by oxysterols via sterol-regulatory element in the promotor, as is found in APOE. Target for cholesterol-lowering therapies: prevastatin, "statins" (Bocan <i>et al.</i> (1998) <i>Atherosclerosis</i> 139:21-30; Farnier <i>et al.</i> (1998) <i>Am J Cardiol</i> 82:3J-10J)
	very low density lipoprotein receptor (VLDLR)	Role in triglyceride metabolism. Marked induction of VLDLR expression observed in fatty streaks and plaques in rabbit atherosclerosis models (Hiltunen <i>et al.</i> (1998) <i>Circulation</i> 97:1079-1086)
20	Microsomal triglyceride transfer protein (MTP)	Catalyses transport of triglyceride, cholesterol ester and phospholipid between phospholipid surfaces. Mutations cause abetalipoproteinemia. Linkage found between MTP genotype and plasma triglyceride levels in a quantitative sib-pair analysis of female dizygotic twins. Inhibitors of MTP normalise atherogenic lipoprotein profiles in an atherosclerotic rabbit model (Wetterau <i>et al.</i> (1992) <i>Science</i> 258:999-1001; Austin <i>et al.</i> (1998) <i>Am J Hum Genet</i> 62:406-419; Wetterau <i>et al.</i> (1998) <i>Science</i> 282:751-754)
	perilipin (PLIN)	Lipid storage droplets of steroidogenic cells are surrounded by perilipins, family of phosphorylated proteins encoded by a single gene, detected in adipocytes and steroidogenic cells. Possible role in lipid metabolism (Brasaemle <i>et al.</i> (1997) <i>J Biol Chem</i> 272:9378-9387)

endothelin-1 (EDN1)	Secretion of EDN1 coincides with the location of native and oxidised low density lipoproteins and occurs in a specific fashion suggesting that EDN1 may be involved in pathophysiological processes such as atherogenesis. Quantitative and qualitative immunohistochemical analysis of anti EDN1 antibodies in the wall layers of human arteries ex vivo suggest that EDN1 is normally expressed exclusively in endothelial cells. However, in cases of coronary artery disease and atherosclerosis, EDN1 expression is enhanced and can be found in the tunica media and vascular smooth muscle cells. Analysis of recombinant EDN1 expression <i>in vitro</i> suggests it influences vascular smooth muscle cell proliferation. Potent vasoconstriction properties (Unoki <i>et al.</i> (1999) Cell Tissue Res 295:89-99; Rossi <i>et al.</i> (1999) Circulation 99:1147-1155; Yoshizumi <i>et al.</i> (1998) Br J Pharmacol 125:1019-1027; Alberts <i>et al.</i> (1994) J Biol Chem 269: 10112-10118)
5 endothelin receptor A (EDNRA)	Mediates action of endothelin1 on vascular smooth muscle migration, proliferation and monocyte/endothelial cell interaction during initiation and progression of atherosclerotic lesion development (Kohno <i>et al.</i> (1998) J Cardiovasc Pharmacol 31:S84-9; Alberts <i>et al.</i> (1994) J Biol Chem 269:10112-10118)
interleukin 6 (IL6)	Inflammatory cytokine present in arterial atherosclerotic wall which is upregulated by platelets to stimulate smooth muscle cell growth. Increased expression of IL6 in atherosclerotic aortas of APOE knockout vs aortas from aged-matched controls. Secretion levels of IL6 is positively associated with increased lesion surface area in APOE aortic tissue samples (Sukovich <i>et al.</i> (1998) Arterioscler Thromb Vasc Biol 18:1498-1505; Loppnow <i>et al.</i> (1998) Blood 91:134-141)
interleukin 1 (IL1)	May contribute to regulation of local pathogenesis in the vessel wall by activation of the cytokine regulatory network. IL-1 antagonist inhibits platelet-induced cytokine production of smooth muscle cells (Loppnow <i>et al.</i> (1998) Blood 91: 134-141)
10 complement protein C8 alpha (C8A)	Complement activation of C8 shown to be an initial event in atherogenesis (Torzewski <i>et al.</i> (1996) Arterioscler Thromb Vasc Biol 16:673-677)
complement component C9 (C9)	Complement activation of C9 shown to be an initial event in atherogenesis (Torzewski <i>et al.</i> (1996) Arterioscler Thromb Vasc Biol 16:673-677)
15 Prostaglandin D2 synthase (PTGDS)	Catalyses conversion of PGH2 to PGD2, a prostaglandin important in smooth muscle contraction/relaxation and potent inhibitor of platelet aggregation. Northern analysis shows strong specific expression in heart. Immunocytochemical localisation to myocardial and atrio endocardial cells, and accumulates in end-stage atherosclerotic plaques. High plasma levels detected in severe angina patients (Eguchi <i>et al.</i> (1997) Proc Natl Acad Sci 94:14689-14694)
Annexin II/lipocortinII (ANX2)	Inhibits phospholipase A2 activity and hence the production of arachidonic acid, the precursor of the inflammatory mediators prostaglandins and leukotrienes. ANX2 is an important anti-inflammatory molecule. Independently binds plasminogen and t-PA and therefore suspected of having a role in atherogenesis. Binding of plasminogen to ANX2 is specifically inhibited by the excess atherogenic Lp(a) (Hajjar <i>et al.</i> (1998) J Investig Med 46(8): 364-369)

	Annexin I/lipocortin (ANXI)	Inhibits phospholipase A2 activity and hence the production of arachidonic acid, the precursor of the inflammatory mediators prostaglandins and leukotrienes. ANXI is an important anti-inflammatory molecule (Wallner <i>et al.</i> (1986) Nature 320:77-81)
5	Prostaglandin- endoperoxide Synthase 2 (PTGS2)	Major mechanism for the regulation of prostaglandin synthesis. Arachidonic acid pathway. Role in inflammation and endothelial cell migration/angiogenesis. Regulated enzyme - major mediator of inflammation. Antiinflammatory glucocorticoids are potent inhibitors of this cyclooxygenase. Over expression of PTGS2 <i>in vitro</i> in rabbit epithelial cells causes increased adhesion to extracellular matrix proteins and inhibition of apoptosis, hallmarks of atherosclerotic plaque formation (Morham <i>et al.</i> (1995) Cell 83:473-482; O'Banion <i>et al.</i> (1992) Proc Natl Acad Sci 89:4888-4892; Tsujii <i>et al.</i> (1995) Cell 83:493-501)
10	insulin-like growth factor binding protein-1 (IGFBP-1)	A study of 218 individuals indicates free IGFBP1 levels are associated with high HDL cholesterol and more favourable cardiovascular outcome. The IGF1/IGFBP1 system found to be associated with cardiovascular risk and atherosclerosis (Janssen <i>et al.</i> (1998) Arterioscler Thromb Vasc Biol 18:277-282)
	Secreted protein, acidic and rich in cysteine (SPARC)	Extracellular glycoprotein secreted by endothelial cells which has a suspected role in calcification of atherosclerotic plaques. Interacts with PDGF-B containing dimers and inhibits binding to its receptors. Expression of SPARC and PDGF is minimal in most adult tissues, but is enhanced following injury and advanced atherosclerotic lesions. Selective expression of SPARC causes rounding of adherent endothelial cells and influences extravasation of macromolecules (Raines <i>et al.</i> (1992) Proc Natl Acad Sci 89:1281-1285; Goldblum <i>et al.</i> (1994) Proc Natl Acad Sci 91:3448-3452)
15	Human NF- kappa-B transcription factor (NFkB)	Activated NF kappa B occurs in atherosclerotic lesions, and regulates the expression of gene important in recruitment of monocytes and inflammatory response. Responsible for cytokine production by smooth muscle cells during atherogenesis (Navab <i>et al.</i> (1995) Am J Cardiol 76:18C-23C; Hernandez-Presa <i>et al.</i> (1998) Am J Pathol 153:1825-1837; Thurberg <i>et al.</i> (1998) Curr Opin Lipidol 9:387-396; Brand <i>et al.</i> (1997) Arterioscler Thromb Vasc Biol 17:1901-1909)
20	angiotensinogen (AGT)	Concentration of angiotensinogen influences the renin-angiotensin system(RAS). Hypertensive mice carrying renin and angiotensinogen transgenes found to have higher total cholesterol levels on an atherogenic diet than their wt counterparts, and atherogenic lesions were 4x larger in surface area. Suggests hypertension induced by activated RAS is important atherogenic factor (Sugiyama <i>et al.</i> (1997) Lab Invest 76:835-842)
	Nitric Oxide Synthase 3 (NOS3)	Mediates basal vasodilation. Regulates the production of nitric oxide, an important signal transduction component and scavenger of reactive oxygen species. Activity of NOS3 appears to be a factor in endothelin/endothelin receptor B mediated endothelial cell migration and angiogenesis. Polymorphism associated with smoking dependent coronary artery disease (Goligorsky <i>et al.</i> (1999) Clin Exp Pharmacol Physiol 26:269-271; Stroes <i>et al.</i> (1998) J Cardiovasc Pharmacol 32:S14-21; Sobue <i>et al.</i> (1998) Horm Res 50:15-24)

Nitric Oxide Synthase 2 (NOS2)	Mediates basal vasodilation. Regulates the production of nitric oxide, an important signal transduction component and scavenger of reactive oxygen species. NOS2, known as inducible NOS is expressed in most cells only after induction by immunologic and inflammatory stimuli, and is upregulated in pathological conditions such as atherosclerosis (Dusting <i>et al.</i> (1998) Clin Expt Pharmacol Physiol 25:S34-41)
--------------------------------	--

5 From a total of 45,233 assembled gene sequences, 34 novel genes were identified, SEQ ID NOs:1-34, that show strong association with 66 known atherosclerosis-associated genes. Initially, the degree of association was measured by probability values using a cutoff p value less than 0.00001. The sequences were further examined to ensure that the genes that passed the probability test had strong association with known atherosclerosis-associated genes. Details of the co-expression patterns for the

10 66 known and 34 novel atherosclerosis-associated polynucleotides are presented in Table 4. The entries in Table 4 are the negative log of the p-value ($-\log p$) for the coexpression of the two genes. The novel atherosclerosis-associated polynucleotides identified are listed in the table by their SEQ ID NOs numbers, and the known genes, by their names or the abbreviations shown in Table 3.

V Novel Genes Associated with Atherosclerosis

15 Using the co-expression analysis method, 34 novel atherosclerosis-associated polynucleotides were identified, SEQ ID NOs:1-34, that exhibit strong association, or co-expression, with 66 known atherosclerosis-associated genes.

Polynucleotides comprising the consensus sequences of SEQ ID NO:1-34 of the present invention were first identified from Incyte bins and assembled as described in Example III. BLAST

20 and other motif searches were performed for SEQ ID NOs:1-34 according to Example VI. The full length and 5'-complete sequences were translated and sequence identity was sought with known sequences.

SEQ ID NO:35 of the present invention was encoded by the nucleic acids of SEQ ID NO:11. SEQ ID NO:35 has 366 amino acids which are encoded by SEQ ID NO:11. Motif analyses of SEQ ID

25 NO:35 shows one potential cAMP- and cGMP-dependent protein kinase phosphorylation sites at residue S343, two potential casein kinase II phosphorylation sites at residues S179 and T351, and four potential protein kinase C phosphorylation sites at residues T29, S85, T269, and T324. Additionally, SEQ ID NO:35 contains a potential sugar transport protein signature sequence from residues L201 to S217.

30 VI Homology Searching for Atherosclerosis-Associated Polynucleotides and Polypeptides

The polynucleotide sequences, SEQ ID NO:1-34, and polypeptide sequence, SEQ ID NO:35, were queried against databases derived from sources such as GenBank and SwissProt. These databases, which contain previously identified and annotated sequences, were searched for regions of similarity using BLAST (Altschul, *supra*). BLAST searched for matches and reported only those that satisfied the

probability thresholds of 10^{-25} or less for nucleotide sequences and 10^{-8} or less for polypeptide sequences.

The polypeptide sequence was also analyzed for known motif patterns using MOTIFS, SPSCAN, BLIMPS, and HMM-based protocols. MOTIFS (Genetics Computer Group, Madison WI) searches polypeptide sequences for patterns that match those defined in the Prosite Dictionary of Protein Sites and Patterns (Bairoch, supra) and displays the patterns found and their corresponding literature abstracts. SPSCAN (Genetics Computer Group) searches for potential signal peptide sequences using a weighted matrix method (Nielsen et al. (1997) Prot Eng 10:1-6). Hits with a score of 5 or greater were considered. BLIMPS uses a weighted matrix analysis algorithm to search for sequence similarity between the polypeptide sequences and those contained in BLOCKS, a database consisting of short amino acid segments, or blocks of 3-60 amino acids in length, compiled from the PROSITE database (Henikoff; supra; Bairoch, supra), and those in PRINTS, a protein fingerprint database based on non-redundant sequences obtained from sources such as SwissProt, GenBank, PIR, and NRL-3D (Attwood et al. (1997) J Chem Inf Comput Sci 37:417-424). For the purposes of the present invention, the BLIMPS searches reported matches with a cutoff score of 1000 or greater and a cutoff probability value of 1.0×10^{-3} . HMM-based protocols were based on a probabilistic approach and searched for consensus primary structures of gene families in the protein sequences (Eddy, supra; Sonnhammer, supra). More than 500 known protein families with cutoff scores ranging from 10 to 50 bits were selected for use in this invention.

VII Labeling of Probes and Hybridization Analyses

Substrate Preparation

Nucleic acids are isolated from a biological source and applied to a substrate for standard hybridization protocols by one of the following methods. A mixture of target nucleic acids, a restriction digest of genomic DNA, is fractionated by electrophoresis through an 0.7% agarose gel in 1xTAE [Tris-acetate-ethylenediamine tetraacetic acid (EDTA)] running buffer and transferred to a nylon membrane by capillary transfer using 20x saline sodium citrate (SSC). Alternatively, the target nucleic acids are individually ligated to a vector and inserted into bacterial host cells to form a library. Target nucleic acids are arranged on a substrate by one of the following methods. In the first method, bacterial cells containing individual clones are robotically picked and arranged on a nylon membrane. The membrane is placed on bacterial growth medium, LB agar containing carbenicillin, and incubated at 37°C for 16 hours. Bacterial colonies are denatured, neutralized, and digested with proteinase K. Nylon membranes are exposed to UV irradiation in a STRATALINKER UV-crosslinker (Stratagene) to cross-link DNA to the membrane.

In the second method, target nucleic acids are amplified from bacterial vectors by thirty cycles of PCR using primers complementary to vector sequences flanking the insert. Amplified target nucleic

acids are purified using SEPHACRYL-400 beads (APB). Purified target nucleic acids are robotically arrayed onto a glass microscope slide (Corning Science Products, Corning NY). The slide is previously coated with 0.05% aminopropyl silane (Sigma-Aldrich) and cured at 110°C. The arrayed glass slide (microarray) is exposed to UV irradiation in a STRATALINKER UV-crosslinker (Stratagene).

5 Probe Preparation

cDNA probes are made from mRNA templates. Five micrograms of mRNA is mixed with 1 µg random primer (Life Technologies), incubated at 70°C for 10 minutes, and lyophilized. The lyophilized sample is resuspended in 50 µl of 1x first strand buffer (cDNA Synthesis systems; Life Technologies) containing a dNTP mix, [α -³²P]dCTP, dithiothreitol, and MMLV reverse transcriptase
10 (Stratagene), and incubated at 42°C for 1-2 hours. After incubation, the probe is diluted with 42 µl dH₂O, heated to 95°C for 3 minutes, and cooled on ice. mRNA in the probe is removed by alkaline degradation. The probe is neutralized, and degraded mRNA and unincorporated nucleotides are removed using a PROBEQUANT G-50 microcolumn (APB). Probes can be labeled with fluorescent markers, Cy3-dCTP or Cy5-dCTP (APB), in place of the radionucleotide, [³²P]dCTP.

15 Hybridization

Hybridization is carried out at 65°C in a hybridization buffer containing 0.5 M sodium phosphate (pH 7.2), 7% SDS, and 1 mM EDTA. After the substrate is incubated in hybridization buffer at 65°C for at least 2 hours, the buffer is replaced with 10 ml of fresh buffer containing the probes. After incubation at 65°C for 18 hours, the hybridization buffer is removed, and the substrate is
20 washed sequentially under increasingly stringent conditions, up to 40 mM sodium phosphate, 1% SDS, 1 mM EDTA at 65°C. To detect signal produced by a radiolabeled probe hybridized on a membrane, the substrate is exposed to a PHOSPHORIMAGER cassette (APB), and the image is analyzed using IMAGEQUANT data analysis software (APB). To detect signals produced by a fluorescent probe hybridized on a microarray, the substrate is examined by confocal laser microscopy, and images are
25 collected and analyzed using GEMTOOLS gene expression analysis software (Incyte Genomics).

VIII Complementary Polynucleotides

Molecules complementary to the polynucleotide, or a fragment thereof, are used to detect, decrease, or inhibit gene expression. Although use of oligonucleotides comprising from about 18 to about 60 base pairs is described, the same procedure is used with larger or smaller fragments or their
30 derivatives (PNAs). Oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and SEQ ID NO:1-34 or fragments thereof. To inhibit transcription by preventing promoter binding, a complementary oligonucleotide is designed to bind to the most unique 5' sequence, most preferably about 10 nucleotides before the initiation codon of the open reading frame. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the mRNA encoding the
35 polypeptide.

IX Producti n of Specific Antibodies

The polypeptides encoded by SEQ ID NO:1-34, or portions thereof, substantially purified using polyacrylamide gel electrophoresis or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols as described in Pound (supra).

5 Alternatively, the amino acid sequence is analyzed using LASERGENE software (DNASTAR, Madison WI) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. Typically, oligopeptides 15 residues in length are synthesized using an ABI 431A Peptide
10 synthesizer (PE Biosystems) using fmoc-chemistry and coupled to keyhole limpet hemocyanin (KLH, Sigma-Aldrich) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (Ausubel, supra) to increase immunogenicity. Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide activity by, for example, binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with
15 radio-iodinated goat anti-rabbit IgG.

X Screening Molecules for Specific Binding with the Polynucleotide or Polypeptide

The polynucleotide, or fragments thereof, or the polypeptide, or portions thereof, are labeled with ³²P-dCTP, Cy3-dCTP, or Cy5-dCTP (APB), or with BIODIPY or FITC (Molecular Probes, Eugene OR), respectively. Libraries of candidate molecules previously arranged on a substrate are
20 incubated in the presence of labeled polynucleotide or polypeptide. After incubation under conditions for either a nucleic acid or amino acid sequence, the substrate is washed, and any position on the substrate retaining label, which indicates specific binding or complex formation, is assayed, and the binding molecule is identified. Data obtained using different concentrations of the polynucleotide or polypeptide are used to calculate affinity between the labeled nucleic acid or protein and the bound
25 molecule.

What is claimed is:

1. A composition comprising an isolated polynucleotide that is coexpressed with one or more known atherosclerosis-associated genes in a plurality of samples and that is selected from the group consisting of:
 - 5 (a) a nucleic acid sequence selected from SEQ ID NOs:1-34;
 - (b) a nucleic acid sequence encoding SEQ ID NO:35;
 - (c) a nucleic acid sequence which is the complement of (a) or (b).
2. A polynucleotide comprising the nucleic acid sequence of SEQ ID NO:8 or the complement thereof.
- 10 3. A composition comprising the polynucleotide of claim 1.
4. A method of using a polynucleotide to screen a library of molecules or compounds to identify at least one ligand which specifically binds the polynucleotide, the method comprising:
 - (a) combining the polynucleotide of claim 1 with a library of molecules or compounds under conditions to allow specific binding, and
 - 15 (b) detecting specific binding, thereby identifying a ligand which specifically binds the polynucleotide.
5. The method of claim 4 wherein the library is selected from DNA molecules, RNA molecules, PNAs, mimetics, and proteins.
6. A ligand identified by the method of claim 4 which modulates the activity of the
20 polynucleotide.
7. A method of using a polynucleotide of to purify a ligand which specifically binds the polynucleotide, the method comprising:
 - (a) combining the polynucleotide of claim 1 with a sample under conditions to allow specific binding,
 - 25 (b) detecting specific binding between the polynucleotide and a ligand,
 - (c) recovering the bound polynucleotide, and
 - (d) separating the polynucleotide from the ligand, thereby obtaining purified ligand.
8. A method for diagnosing a disease or condition associated with the altered expression of a polynucleotide that is coexpressed with one or more known atherosclerosis-associated genes in a
30 sample, the method comprising the steps of:
 - (a) hybridizing the composition of claim 1 to a sample under conditions to form one or more hybridization complexes;
 - (b) detecting the hybridization complexes; and
 - (c) comparing the levels of the hybridization complexes with the level of hybridization
35 complexes in a non-diseased sample, wherein the altered level of hybridization complexes compared

with the level of hybridization complexes of a non-diseased sample indicates the presence of the disease or condition.

9. An expression vector comprising the polynucleotide of claim 2.

10. A host cell comprising the expression vector of claim 9.

5 11. A method for producing the polypeptide, the method comprising:

(a) culturing the host cell of claim 10 under conditions for expression of the polypeptide,

(b) recovering the polypeptide from cell culture.

12. A substantially purified polypeptide comprising the product of a gene that is coexpressed with one or more known atherosclerosis-associated genes in a plurality of samples.

10 13. The polypeptide of claim 12, comprising a polypeptide sequence selected from

(a) the polypeptides encoded by SEQ ID NOs:1-34; and

(b) an oligopeptide sequence comprising at least 6 sequential amino acids of the polypeptide sequence of a).

14. The polypeptide comprising the amino acid sequence of SEQ ID NO:35.

15 15. A pharmaceutical composition comprising a polypeptide of claim 12 and a pharmaceutical carrier.

16. A method for using a polypeptide to screen a library of molecules or compounds to identify at least one ligand which specifically binds the polypeptide, the method comprising:

(a) combining the polypeptide of claim 12 with the library of molecules or compounds

20 under conditions to allow specific binding, and

(b) detecting specific binding between the polypeptide and ligand, thereby identifying a ligand which specifically binds the polypeptide.

17. The method of claim 16 wherein the library is selected from DNA molecules, RNA molecules, PNAs, mimetics, proteins, agonists, antagonists, and antibodies.

25 18. A ligand identified by the method of claim 16 which modulates the activity of the polypeptide.

19. A method of using the polypeptide to purify a ligand from a sample, the method comprising:

30 (a) combining the polypeptide of claim 12 with a sample under conditions to allow specific binding,

(b) detecting specific binding between the polypeptide and a ligand,

(c) recovering the bound polypeptide, and

(d) separating the polypeptide from the ligand, thereby obtaining purified ligand.

20. A method for treating a disease associated with the altered expression of a gene that is
35 coexpressed with one or more known atherosclerosis-associated genes in a subject in need, the method

comprising the step of administering to the subject in need the pharmaceutical composition of claim 15 in an amount effective for treating the disease.

SEQUENCE LISTING

<110> INCYTE GENOMICS, INC.
JONES, Karen Anne
VOLKMUTH, Wayne
WALKER, Michael

<120> ATHEROSCLEROSIS-ASSOCIATED GENES

<130> PB-0013 PCT

<140> To Be Assigned

<141> Herewith

<150> 09/349,015

<151> 1999-07-07

<160> 35

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1334

<212> DNA

<213> HOMO SAPIENS

<222> 674, 735, 788

<400> 1

aggcctccct	ccacctgtct	tctcagagca	gataatggca	agcatggctg	ccgtgctcac	60
ctgggctctg	gctcttcttt	cagcgthttc	ggccaccag	gcacggaaag	gcttctggga	120
ctacttcagc	cagaccagcg	gggacaaagg	cagggtggag	cagatccatc	agcagaagat	180
ggctcgcgag	cccgcgaccc	tgaaagacag	ccttgagcaa	gacctcaaca	atatgaacaa	240
gttctctggaa	aagctgaggc	ctctgagtgg	gagcgaggct	cctcggctcc	cacaggaccc	300
ggtgggcatg	cggcggcagc	tgcaggagga	gttgaggagg	gtgaaggctc	gcctccagcc	360
ctacatggca	gaggcgacag	agctgggtgg	ctggaatttg	gagggcttgc	ggcagcaact	420
gaagccctac	acgatggatc	tgatggagca	ggtggccctg	cgctgtcagg	agctgcagga	480
gcagttgcg	gtggtggggg	aagacaccaa	ggcccagttg	ctggggggcg	tggacgaggc	540
ttgggtcttg	ctgcaggggac	tgacagagccg	cgtggtgcac	cacaccggcc	gcttcaaaga	600
gctcttccac	ccatacgccg	agagcctggt	gagcggcatc	gggcgccacg	tgcaggagct	660
gcaccgcagt	gtgntccgca	cgcccccgcc	agccccgcgc	gcctcagtcg	ctgcgtgcag	720
gtgctctccc	ggaantcagc	ctcaaggcca	aggccctgca	cgcacgcac	cagcagaacc	780
tggaccantg	cgccaagagc	tcagcagagc	ctttgcaggc	actgggactg	aggaaggggc	840
cgccccggac	ccccagatgc	tctccgagga	ggtgcgccag	cgacttcagg	ctttccgcca	900
ggacacctac	ctgcagatag	ctgccttcac	tcgcgccatc	gaccaggaga	ctgaggaggt	960
ccagcagcag	ctggcgccac	ctccaccagg	ccacagtgcc	ttcgccccag	agtttcaaca	1020
aacagacagt	ggcaagggtc	tgagcaagct	gcaggcccgt	ctggatgacc	tgtgggaaga	1080
catcactcac	agccttcatg	accagggcca	cagccgtctg	ggggaccctc	gaggatctac	1140
ctgcccaggc	ccattcccag	cttcttgtct	ggggagccct	ggctctgagc	ctctagcatg	1200
gttcagtcct	tgaaagtggc	ctgttgggtg	gaggggtggaa	ggctcctgtg	aggacagggg	1260
ggccaccaa	ggggctgctg	tctcctgcac	atccagcctc	ctgcgactcc	ccaatgcagg	1320
atgcattcat	tcac					1334

<210> 2

<211> 1702

<212> DNA

<213> HOMO SAPIENS

<400> 2

cggtccact	gcaccctgga	gaacgagcct	ttgcggggtt	tctcctggct	gtcctccgac	60
cccggcggtc	tcgaaagcga	cacgctgcag	tgggtggagg	agccccaacg	ctcctgcacc	120

gcgcggagat	gcgcggtact	ccaggccacc	gggtggggtcg	agcccgcagg	ctggaaggag	180
atgcgatgcc	acctgcgcgc	caacggctac	ctgtgcaagt	accagtttga	ggtcttgtgt	240
cctgcgccgc	gccccggggc	cgctctaac	ttgagctatc	gcgcgccctt	ccagctgcac	300
agcgcgcctc	tggacttcag	tccacctggg	accgaggtga	gtgcgctctg	ccgggggacag	360
ctccccgatct	cagttacttg	catcgcggac	gaaatcggcg	ctcgtctggga	caaactctcg	420
ggcgatgtgt	tgtgtccctg	ccccggggagg	tacctccgtg	ctggcaaagt	cgcagagctc	480
cctaactgcc	tagacgactt	gggaggcttt	gcctgcgaat	gtgctacggg	cttcgagctg	540
gggaaggacg	gccgctcttg	tgtgaccagt	ggggaaggac	agccgacctt	tgggggggacc	600
ggggtgcccc	ccaggcgccc	gccggccact	gcaaccagcc	ccgtgccgca	gagaacatgg	660
ccaatcaggg	tcgacgagaa	gctgggagag	acaccacttg	tccctgaaca	agacaattca	720
gtaacatcta	ttcctgagat	tcctcgatgg	ggatcacaga	gcacgatgtc	tacccttcaa	780
atgtcccttc	aagccgagtc	aaaggccact	atcaccccat	cagggagcgt	gatttccaag	840
tttaattcta	cgacttcctc	tgccactcct	caggctttcg	actcctcctc	tgccgtgggtc	900
ttcatatttg	tgagcacagc	agtagtagtg	ttgggtgatct	tgaccatgac	agtactgggg	960
cttgtcaagc	tctgctttca	cgaaagcccc	tcttcccagc	caaggaagga	gtctatgggc	1020
ccgcggggcc	tggagagtga	tcctgagccc	gctgcttttg	gctccagttc	tgacacattgc	1080
acaaacaatg	gggtgaaaagt	cggggactgt	gatctgcggg	acagagcaga	gggtgccttg	1140
ctggcggagt	cccctcttgg	ctctagtgat	gcatagggaa	acaggggaca	tgggcactcc	1200
tgtgaacagt	ttttcacttt	tgatgaaacg	gggaaccaag	aggaacttac	ttgtgtaact	1260
gacaatttct	gcagaaatcc	cccttcctct	aaattccctt	tactccactg	aggagctaaa	1320
tcagaactgc	acactccttc	cctgatgata	gaggaagtgg	aagtgccttt	aggatgggtga	1380
tactggggga	ccgggtagtg	ctggggagag	atattttctt	atgtttattc	ggagaatttg	1440
gagaagtgat	tgaacttttc	aagacatttg	aaacaaatag	aacacaatat	aatttacatt	1500
aaaaaataat	ttctaccaa	atggaaaagga	aatgttctat	gttgttcagg	ctaggagtat	1560
attggttcga	aatcccaggg	aaaaaaaataa	aaataaaaaa	ttaaaggatt	gttgataaaa	1620
aaaaaaaaaa	aaaaagatct	ttaattaagc	ggcccaagct	tattcccttt	agtgagggtt	1680
aatttttagct	tgcaactggcc	ac				1702

<210> 3
 <211> 586
 <212> DNA
 <213> HOMO SAPIENS
 <222> 48, 66, 560, 574, 577, 580

<400> 3						
tcgaggactc	cgccaactac	agctgcgtct	acgtggacct	gaagccgnct	ttcgggggct	60
acgcgnccag	cgagcgcttg	gagctgcacg	tggacggacc	ccctcccagg	cctcagctcc	120
gggcgacgtg	gagtggggcg	gtcctggcgg	gccgagatgc	cgctcctgcg	tgcgagggac	180
ccatccccga	cgtcaccttc	gagctgctgc	gcgagggcga	gacgaaggcc	gtgaagacgg	240
tccgcacccc	cggggccgcg	gcgaacctcg	agctgatctt	cgtggggccc	cagcacgccg	300
gcaactacag	gtgccgctac	cgctcctggg	tgccccacac	cttcgaatcg	gagctcagcg	360
accctgtgga	gctcctggtg	gcagaaagct	gatgcagccg	cgggcccagg	gtgctgttgg	420
tgctctcaga	agtgcggggg	attctggact	ggctccctcc	cctcctgttg	cagcacaagg	480
ccgggggtctc	tgggggggctg	gagaagcctc	cctcattcct	cccaggaatt	aataaatgtg	540
aagagagctc	tgttttaaaan	aaaaaaaaag	aaanaanaan	aaccaa		586

<210> 4
 <211> 433
 <212> DNA
 <213> HOMO SAPIENS

<400> 4						
ctcaagaccc	agcagtggga	cagccagaca	gacggcacga	tggcactgag	ctcccagatc	60
tgggcgcgtt	gcctcctgct	cctcctcctc	ctcgccagcc	tgaccagtgg	ctctgttttc	120
ccacaacaga	cgggacaact	tgacagagctg	caaccccagg	acagagctgg	agccagggcc	180
agctggatgc	ccatgtttcca	gaggcgaagg	aggcgagaca	cccacttccc	catctgcatt	240
ttctgctgcy	gctgctgtca	tcgatcaaag	tgtgggatgt	gctgcaagac	gtagaacctc	300
cctgccttgc	ccccgtcccc	tccttctcct	atatttctct	gctgccccag	aacataggtc	360
ttggaataaa	atggctggtt	cttttgtttt	ccaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	420
aaaaaaaaaa	aaa					433

<210> 5
 <211> 752

<212> DNA
<213> HOMO SAPIENS

<400> 5

attgtacact	ttaaaataat	ggaattttac	agtaagtga	gtatgtatcg	atgaagctat	60
taaaacatct	atattattagc	tcaaattttct	acaggtcaga	attctggcat	ggagtggctg	120
cattctttgt	ttaagctgaa	atcaagggtgt	tgtttgccg	tggtctcacc	tgaagctcag	180
agttcacttc	caagctcatt	tttgtcctta	gcagaattga	gtgtcttgca	attgtagaac	240
tgaggtcttg	gcttgtctgc	tgtcagcagg	ggactgctcc	ctgcttctag	aggccaccgg	300
tttccctcgt	tatgtggccc	cttccatttt	caggccagca	ataatgtgtt	gaatacttcc	360
tatgcttcaa	atctctggct	tctgctacca	gctggagaaa	aaactctctg	cttgtagagg	420
gctcatgtga	tttacttagt	ctttgtctta	aggtcaattt	atttgggtact	tgggatttta	480
attgtatctg	tatgtttcca	tcaaggcaat	aactgtatta	gtgtttgaat	aaataaccag	540
gtaatctggg	aattttaccat	actggtaatc	tgacagggag	atgggaattc	atctttataa	600
ttctgcttac	cacaaaccat	gtctgtgctt	attttctttg	gggaagagtt	gtctgtgact	660
gtcacttagt	ttgaggttcc	atgttgctga	gattctgtcc	agtattttga	cctcttcccc	720
aaatctgggc	ttcagaacca	tctcttagga	gc			752

<210> 6
<211> 944
<212> DNA
<213> HOMO SAPIENS

<400> 6

tcttggggccg	agaattttttt	ttttttttttt	tttgettggt	cgggtaattt	tcattccaaa	60
taaacttatc	acaaaaaaac	tcagcttccc	aaggctcatc	ccccgctgcc	agatacatatc	120
ttatctctga	aagagtttgg	aagatggacc	tttcaattcc	tctacaatta	gtagctgagt	180
tacagagtaa	cctgccagca	atcctatcag	cattcatcag	actattttaa	tagagcaaaag	240
tccacaaaaa	gttccactga	gacatgctga	gcaaaggccg	gagccccaga	agaaaacaag	300
tacagactca	gaggaaagct	gccctggctc	tgagtgtgac	tcccatgggc	cccgtggggg	360
ctgtgtgggt	ggcaatgagc	tctgtgctgt	cagctttcat	gagggagctc	cctggctggg	420
tcctgttctt	tgggggtctt	ctccccatga	ctttgtctgt	gtctctctc	atcgccctact	480
tcaggatcaa	actgattgag	gttaatgaag	aactgtccca	gaactgtgat	cgccaacata	540
atcccaagga	tggctcttcc	ctgtaccaga	gaatgaaatg	gacgtgaagt	tgggtgacttt	600
ccaataacta	aagcacaatg	agtttctact	ggtcagcaag	caatggccaa	cagttcagct	660
aataaagtag	ggtgataaac	tagaaccata	gcaaaataga	aagaatacta	agatactcat	720
tctgaaccat	actgaaaagt	ggcagctatt	atctaagggg	acttctcaga	gactcagtat	780
aacagcagct	cttgaaaagt	accaagaatg	gatttctctg	gtatatacac	tggacacatt	840
gtaacttttt	aactttttatt	gtgactgtgt	ctgctctaaa	cggcatattt	aaaaaataaa	900
attctgcagc	atcttactac	ataaaaaaaaa	aaaaaaaaaa	aaaa		944

<210> 7
<211> 868
<212> DNA
<213> HOMO SAPIENS

<400> 7

cctccctccg	cgagctggac	gctccgcagc	ccgcccgcga	gccggcccgc	cgccgcgcgc	60
aggaatccct	ggataaagac	cagctcaacc	atcgctgaga	aaacagacct	aggcttccca	120
gggcggttaa	cccgcgggcc	tctgggcaga	gactaaaaga	caaaacaaaa	taaaacaaca	180
acaaaaaact	cccagtggtg	ttcctactct	tctttgtctt	ggaggaaagc	aaagggagag	240
aaatggactt	caccagtggg	ctttggcttc	atcaattcac	aggaaatggc	atcaagatgg	300
ttcaactaag	acatgatcac	taaaaacatt	ataataatac	ctttttgaaa	aactcagttt	360
ctcctgttta	ctaaatattt	atttcatcaa	catgggctgc	gttccactgt	gtcaggattc	420
tgcatgtggg	tggagcactg	ttccagcctg	agaagatggg	tctgaggcca	cttagcaaga	480
cattttccag	catgagcagg	tttctctgtg	gaaatagtga	cacctgttct	ggtgtgttgt	540
ctttcctcag	ggaacttaag	gggtacaaaag	ctcctgaaaa	tggtctttat	gctgggttgaa	600
gctcttatgt	cgctgtactg	attccctacg	atgcagattt	gaatcacaga	gtaattaaaa	660
tatggatcaa	ataaggctgg	ggctcaccaa	ggctgaaagc	tgtagccatt	caaggcatca	720
tttctgtcat	gaaaatatag	gaccttttca	aaacatgcct	tcaggaaggt	gttctctttt	780
caaacaaaag	tctaagtact	gcataactct	tcttgaccac	atcttacact	ttctctagac	840
ttgcttattt	acagctactg	gaacaaaa				868

<210> 8
 <211> 3111
 <212> DNA
 <213> HOMO SAPIENS
 <222> 44

<400> 8
 cgagggcgga cgcaaagaac ggggaggacc tctgggtgcc tgcnggggag ctgctccagc 60
 cgggcccgcg ggagcgggtg ggagagcacc gcgcagccgc ccctccacgc gcccgcgccag 120
 ccgcgttcgc ccactgggct ctcccggctg cagtgccagg gcgcaggacg cggccgatct 180
 cccgctcccc ccacctccgc caccatgctg ctccccccagc tctgctggct gccgctgctc 240
 gctgggctgc tcccgcgggt gcccgcctcag aagttctcgg cgctcacgtt ttgagagtg 300
 gatcaagata aagacaagga ttgtagcttg gactgtgcgg gttcgccccca gaaacctctc 360
 tgcgcatctg acggaaggac ctctctttcc cgttgtgaat ttcaacgtgc caagtgcaaa 420
 gatccccagc tagagattgc atatcgagga aactgcaaag acgtgtccag gtgtgtggcc 480
 gaaaggaagt ataccagga gcaagcccgg aaggagtttc agcaagtgtt cattcctgag 540
 tgcaatgacg acggcaccta cagtcaggtc cagtgtcaca gctacacggg atactgctgg 600
 tgcgtcacgc ccaacgggag gcccatcagc ggcactcccg tggcccacaa gacgcccgg 660
 tgcccggggt ccgtaaatga aaagttaccc caacgcgaag gcacaggaaa aacagctgat 720
 gccgcagctc cagcgttgga gactcagcct caaggagatg aagaagatat tgcatacgt 780
 taccctaccc ttggactga acaggttaaa agtcggcaga acaaaaccaa taagaattca 840
 gtgtcatcct gtgaccaaga gcaccagtct gccctggagg aagccaagca gcccaagaac 900
 gacaatgtgg tgatccctga gtgtgcgcac ggcgccctct acaagccagt gcagtgccac 960
 ccctccacgg ggtactgctg gtgcgtcttg gtggacacgg ggcgccccat tcccggcaca 1020
 tccacaaggt acgagcagcc gaaatgtgac aacacgggccc agggcccacc cagccaaagc 1080
 ccgggacctg tacaagggcc gccagctaca aggttgtccg ggtgccaaaa agcatgagtt 1140
 tctgaccagc gttctggacg cgctgtccac ggacatggtc cacgcgcct cccgcccctc 1200
 ctctcgtca ggcaggctct cagaaccgca cccagccat accctagagg agcgggtggt 1260
 gcaactgtac ttcaaactac tggataaaaa ctccagtga gacatcggca aaaaggaaat 1320
 caaaccttc aagaggttcc ttcgcaaaaa atcaaagccc aaaaaatgtg tgaagaagt 1380
 tgttgaatac tgtgacgtga ataatacaca atccatctcc gtacaagaac tgatgggctg 1440
 cctgggcgtg gcgaaagagg acggcaaaagc ggacaccaag aaacgccaca ccccagagg 1500
 tcatgctgaa agtacgtcta atagacagcc aaggaaacaa ggataaatgg ctcatacccc 1560
 gaaggcagtt cctagacaca tgggaaattt cctcaccaca agagcaatta agaaaacaaa 1620
 aacagaaaca catagtattt gcactttgtg cttaaattgt aaattcactt tgtagaaatg 1680
 agctatttaa acagactgtt ttaatctgtg aaaatggaga gctggcttca gaaaattaat 1740
 cacataccaa tgtatgtgtc ctcttttgac cttggaaatc tgtatgtggt ggagaagtat 1800
 ttgaatgcat ttaggcttaa tttcttcgcc ttccacatgt taacagtaga gctctatgca 1860
 ctccggctgc aatcgtatgg ctttctctaa cccctgcagt cacttcagga tgccgtgct 1920
 tacagcattg tggaaatcat ttggaatcag cacatgtcca tggaaagttg tgatgtacgg 1980
 ccgaccctac aggcagttaa catgcatggg ctggtttgtt tcttgggatt tctgttagt 2040
 ttgtcttgtt ttgctttcca gagatcttgc tcatacaatg aatcacgcaa ccactaaagc 2100
 tatccagtta agtgcaagta gttcccctgg aggaaataat attttcaaac tgtcgttgg 2160
 gtgatacttt ggctcaaagg atctttgctt ttccatttta agcttctgtt ttgagttttg 2220
 ccctggggct tgaatgagtc ccagagagtc gttcggatgg tgggaggctg cctaggaggc 2280
 agtaaatcca gttcacagt cctgggaggg gcccatcctt ccaaaatgta aatccagttc 2340
 gcggtgtgac cgagctgggc taacaggctt gtctgcctgg ttttcctacc tacacgtgga 2400
 cattattctc ctgatacctc tacctggttc caccacagg ctaccggaag gtaaaatctt 2460
 cacctgaacc aattatgagc agtctcctta ctgaaggtag agccggatag gtggtgcccc 2520
 cggggctggt gttggcagcc ggggggaggt gcctgaggg cccacgggt ctttctgct 2580
 tttctgaatg catcaagggt acgagaactt gccaatggga aattcatccg agtggcactg 2640
 gcagagaagg ataggagtgg aatgcccaca cagtgaacaa cagaactgggt ctgcgtgcat 2700
 aaccagctgc caccctcagg cctgggcccc agagctcagg gcacccagtg tcttaaggaa 2760
 ccatttggag gacagtctga gagcaggaac ttcaagctgt gattctatct cggctcagac 2820
 ttttggttgg aaaaagatct tcatggcccc aaatcccctg agacatgcct tgtagaatga 2880
 ttttgtgatg ttgtgatgct tgtggagcat cgcgtaaggc ttcttgctta ttaaaactgt 2940
 gcaaggtaaa aatcaagcct ttggagccac agaaccagct caagtacatg ccaatgttgt 3000
 ttaagaaaca gttatgatcc taaacttttt ggataatctt ttatatctt gacctttgaa 3060
 tttaatcatt gttcttagat taaaataaaa tatgctattg aaactaaaaa a 3111

<210> 9
 <211> 2311
 <212> DNA

<213> HOMO SAPIENS

<222> 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 2288, 2295, 2296, 2297, 2298, 2299

<400> 9

gccgctcgcc	cacggactcc	gacgtgtccc	tgcactccga	ggactccggg	gctaagtctc	60
caggcatcct	gggctacaat	atctgtcccc	gcgggtggaa	tggcagcctt	cggtcaagc	120
gtggcagcct	ccccgccgag	gcctcctgca	ccacctagag	ccccaccccc	gacccccacc	180
cgggagggca	gagccagaag	aaggctcatt	agacctgggg	gacccaaagg	gtctggcctc	240
tttgggcagc	cccagagatg	aggggtcagc	agaggagagc	tctgggggtg	gggatgggtt	300
agggagcga	gcttgagtgc	tagcccttgc	tctcattcag	ctgttggtg	acctgggta	360
agacccttcc	ttgtttgacc	ctcagctttc	ccatctgttt	aatgggtggc	ttggccaagg	420
caatccacaa	acgtcaaaat	tccccttccc	atcagtacac	acaccgatgc	acannnnnnn	480
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	540
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	600
nnntagtta	gtgccttgga	tgaggcgggg	cagtgtgtat	atggaccctt	ggacttgcta	660
ccttcagggt	tccatactcg	tccctcccc	cctggctctg	ctgtctggag	tctggcaagc	720
ggggtgtgtt	cagaagggtc	taggcctgtg	tgcagtgtcc	aggcactggc	ctgaccatcc	780
ggctccctgg	gcaccaagtc	ccagggcagg	agcagctgtt	ttccatccct	tcccagacaa	840
gctctatttt	tatcacaatg	accttttagag	aggtctccca	ggccagctca	aggtgtccca	900
ctatcccttc	tggagggaag	aggcaggaaa	attctccccg	ggtccctgtc	atgctacttt	960
ctccatccca	gttcagactg	tccaggacat	cttatctgca	gccataagag	aattataagg	1020
cagtgtattc	ccttagggcc	aggacttggg	cctccagctc	atctgttcc	tctgggcccc	1080
ttcatggcca	ggttctgggc	tcaaagctga	actggggaga	gaagagatac	agagctacca	1140
tgtgacttta	cctgattgcc	ctcagtttgg	ggttgcttat	tgggaaagag	agagacaaag	1200
agttacttgt	tacgggaaat	atgaaaagca	tggccaggat	gcatagagga	gattctagca	1260
ggggacagga	ttggctcaga	tgacccctga	gggctcttcc	agtcttgaaa	tgcattccat	1320
gatattagga	agtcgggggt	gggtggtggt	ggtgggctag	ttgggcttga	atttaggggc	1380
cgatgagctt	gggtacgtga	gcagggtgtt	aagttagggg	ctgcctgtat	ttctggtccc	1440
cttgggaat	gtccccttct	tcagtgtcag	acctcagtc	cagtgtccat	atcgtgcccc	1500
gaaaagtaga	cattatcctg	ccccatccct	tcccagctgc	actctgacct	agctagtgcc	1560
tgggtgccag	tgacctgggg	gagcctggct	gcaggccctc	actgggtccc	taaaccttgg	1620
tggctgtgat	tcagggtcccc	aggggggact	cagggaggaa	tatggctgag	ttctgtagtt	1680
tccagagttg	ggctggtaga	gctttctaga	ggttcagaat	attagcttca	ggatcagctg	1740
ggggtatgga	attggctgag	gatcaaactg	atgttaggtg	aaggatacca	ggatgttgct	1800
aaaggtgagg	gacagtttgg	gtttgggact	taccgggggtg	atgttagatc	tggaaacccc	1860
aagtgaggct	ggaggagggt	aagggtcagta	tgggaagatag	ggttggggaca	gggtgctttg	1920
gaatgaaaga	gtgaccttag	agggctcctt	gggcctcagg	aatgctcctg	ctgctgtgaa	1980
gatgagaagg	tgctcttact	cagttaatga	tgagtgaacta	tatttaccac	agccccctacc	2040
tgctgctggg	tcccttgtag	cacaggagac	tgggggctaag	ggccccctcc	agggaaggga	2100
caccatcagg	cctctggctg	aggcagtagc	atagaggatc	catttctacc	tgcatttccc	2160
agaggactag	caggaggcag	ccttgagaaa	cgggcagttc	ccaagccagc	gcctggctgt	2220
tctctcattg	tactgccc	ctcccccaac	tctcctctaa	cccactagag	attgcctgtg	2280
tctgcccct	tgccctctgt	agaatgcagc	tctggccctc	aataaatgct	tctgcatc	2311
taaaaanaa	aaaannnnna	aaaaaaaaag	g			

<210> 10

<211> 1866

<212> DNA

<213> HOMO SAPIENS

<400> 10

agcttttgtt	cacactttaa	atagcagtc	cagaatgatt	tactacaga	ctctctggaa	60
agcctgggag	ctgaattccg	gaagatcccc	acatcgatga	aagcaaagcg	aagccaccaa	120
gccatcatca	tgtccacgtc	gctacgagtc	agccccatcca	tccatggcta	ccacttcgac	180
acagcctctc	gtaagaaagc	cgtgggcaac	atctttgaaa	acacagacca	agaatcacta	240
gaaaggctct	tcagaaactc	tgagagacaag	aaagcagagg	agagagccaa	gatcattttt	300
gccatagatc	aagatgtgga	ggagaaaacg	cgtgccctga	tggccttgaa	gaagaggaca	360
aaagacaagc	ttttccagtt	tctgaaactg	cggaaatatt	ccatcaaagt	tactgaaga	420
gaagaggatg	gataaggacg	ttatccaaga	atggacattc	aaagaccaag	tgagtttgtg	480

agattctaac	agatgcagca	ttttgctgct	accttacaag	cttctcttct	gtcaggactc	540
cagaggctgg	aaagggaccg	ggactggaaa	gggaccagga	ctgaacagac	tggttacaaa	600
gactccaaac	aatttcatgc	cctgtgctgt	tacagaggag	aacaaaatgc	tttcagcaag	660
gatttgaaaa	ctcttccgtc	cctgcaggaa	aggattgatg	ctgatagaag	agcctggaca	720
gatgtaatga	gaactaaaga	aaacagatgg	ctggagatga	catttatcca	gggtcacttt	780
gtcaggccct	aggacttaaa	tccaagttga	actttttttt	ttttttaacc	aaatagatag	840
gggaagggag	gagggagagg	gaggacaggg	agagaaaata	ccatgcataa	attgtttact	900
gaatttttat	atctgagtgt	tcaaaatatt	tccaagcctg	agtattgtct	attggatatag	960
atttttagaa	atcaataatt	gattattttat	ttgcacttat	tacaatgcct	gaaaaagtgc	1020
accacatgga	tgtaagtag	aaattcaaga	aagtaagatg	tcttcagcaa	ctcagtaaaa	1080
ccttacgcca	ccttttggtt	tgtaaaaggt	tttttataca	tttcaaacag	gttgccacaaa	1140
agttaaaata	atggggtcct	ttataaatcc	aaagtactgt	gaaaacattt	tacatatattt	1200
ttaaatcttc	tgactaatgc	taaaacgtaa	tctaattaaa	tttcatacag	ttactgcagt	1260
aagcattagg	aagtgaatat	gatatacaaa	atagtttata	aagactctat	agtttctata	1320
atttatttta	ctggcaaatg	tcatgcaaca	ataataaatt	attgtaaact	ttgtggcctt	1380
tggtctgtga	tgcttggtct	caaaggaaaa	aataagatgg	taaatgttga	tatttacaaa	1440
cttttctaaa	gatgtgtctc	taacaataaa	agttaatttt	agagtagttt	tatattaatt	1500
accaaacttt	ttcaaaacaa	attcttacgt	caaatatctg	ggaagtctct	ctgtcccaat	1560
cttaaaatat	aaaatataga	tatagaagtt	catagattga	ctccttgcca	tttctattta	1620
tgtatccatt	aaggatgagt	tttaaaaggg	tttctcttca	tacttttgaa	aaatttcttc	1680
tatgattaca	gtagctatgt	acatgtgtac	atctattttt	cccaagcaat	atgttttggg	1740
tttagagtct	gagtgatgac	caagattctg	tgtgttacta	ctgtttgttt	aataggaaca	1800
aatatagaaa	taatattatc	tctttgctta	tttcccgtaa	aaactataat	aaaatgtttc	1860
taggaa						1866

<210> 11

<211> 1929

<212> DNA

<213> HOMO SAPIENS

<400> 11

gctgcctgcc	ggtgctcttc	gtggctctgg	gcattggcctc	ggaccccatc	ttcacgctgg	60
cgcccccgct	gcattgccac	tacggggcct	tccccctaa	tgcctctggc	tgaggagcagc	120
ctcccaatgc	cagcggcgctc	agcgtcgcca	gcgctgccct	agcagccagc	gccgccagcc	180
gtgtcgccac	cagtaccgac	ccctcggtga	gcggcttcgc	cccgcgggac	ttcaaccatt	240
gccctcaagg	attgggacta	taatggcctt	cctgtgtcca	ccaccaacgc	catcggccag	300
tgggatctgg	tgtgtgacct	gggctggcag	gtgatcctgg	agcagatcct	cttcactctg	360
ggctttgcct	ccggctacct	gttcctgggt	taccccgag	acagatttgg	ccgtcgcggg	420
attgtgctgc	tgaccttggg	gctggtgggc	ccctgtggag	taggaggggc	tgctgcaggc	480
tctccacag	gcgtcatggc	cctccgattc	ctcttgggct	ttctgcttgc	cggtgttgac	540
ctgggtgtct	acctgatgcg	cctggagctg	tgcgacccaa	cccagaggct	tgggtggcc	600
ctggcagggg	agttgggtggg	ggtgggaggg	cacttctgt	tcctgggect	ggccttgtc	660
tctaaggatt	ggcgattcct	acagcgaatg	atcaccgctc	cctgcatect	cttctgttt	720
tatggctggc	ctggtttgtt	cctggagtcc	gcacggtggc	tgatagtga	gcggcagatt	780
gaggaggctc	agtctgtgct	gaggatcctg	gctgagcgaa	accggcccca	tgggcagatg	840
ctgggggagg	aggcccagga	ggccctgcag	gacctggaga	atacctgccc	tctccctgca	900
acatcctcct	tttcccttgc	ttccctcctc	aactaccgca	acatctggaa	aaatctgctt	960
atcctgggct	tcaccaactt	cattgcccct	gccattcgcc	actgctacca	gcctgtggga	1020
ggaggaggga	gcccacgga	cttctacctg	tgctctctgc	tgccagcg	caccgcagcc	1080
ctggcctgtg	tcttccctggg	ggtcacccgtg	gaccgatttg	gccgcggggg	catccttctt	1140
ctctccatga	cccttaccgg	cattgcttcc	cttggtcctgc	tgggctgtg	ggattatctg	1200
aacgaggctg	ccatccacac	ttctctctgc	cttgggctct	tctcctccca	agctgcgcgc	1260
atcctcagca	ccctcccttgc	tgctgaggtc	atccccacca	ctgtccgggg	ccgtggcctg	1320
ggcctgatca	tggtcttagg	ggcgcttgga	ggactgagcg	gcccggccca	gcgcctccac	1380
atggggccatg	gagccttct	gcagcacgtg	gtgctggcgg	cctgcgccct	cctctgcatt	1440
ctcagcatta	tgctgtgcc	ggagaccaag	cgcaagctcc	tgcccagagg	gctccggggac	1500
ggggagctgt	gtcgccggcc	ttccctgctg	cggcagccac	cccctaccgc	ctgtgaccac	1560
gtcccgctgc	ttgcccaccc	caacctggcc	ccttgagcgg	cctctgagta	ccctggcggg	1620
aggctggccc	acacagaaag	gtggcaagaa	gatcggggaag	actgagtagg	gaaggcaggg	1680
ctgcccagaa	gtctcagagg	cacctcacgc	cagccatcgc	ggagagctca	gagggccgtc	1740
cccaccctgc	ctcctccctg	ctgctttgca	ttcacttct	tggccagagt	caggggacag	1800
ggagagagct	ccacactgta	accactgggt	ctgggctcca	tctgcgccc	aaagacatcc	1860
accagacct	cattatttct	tgtcttatca	ttctgtttca	ataaagacat	ttggaataaa	1920

aaaaaaaa

1929

<210> 12
 <211> 1831
 <212> DNA
 <213> HOMO SAPIENS

<400> 12
 ctggagccgc cctgggtgtc agcgggtcgg cteccgcgca cgcctccggcc gtcgcgcacc 60
 tcgggcacct gcagggtccgt ggcgtcccg cggctgggcgc ccctgactcc gteccggcca 120
 gggagggcca tgatttccct cccggggccc ctggtgacca acttgctgcg gtttttgttc 180
 ctggggctga gtgccctcgc gccccctcg cgggcccagc tgcaactgca cttgcccgcc 240
 aaccggttgc aggcgggtgga gggaggggaa gtggtgcttc cagcgtggta caccttgcac 300
 ggggaggtgt cttcatccca gccatgggag gtgccctttg tgatgtggtt cttcaaacag 360
 aaagaaaaggt aggatcagggt gttgtcctac atcaatgggg tcacaacaag caaacctgga 420
 gtatccttgg tctactccat gccctcccg aacctgtccc tgcggctgga gggctctccag 480
 gagaaagact ctggccccta cagctgtctc gtgaatgtgc aagacaaaaca aggcaaatct 540
 agggggccaca gcatcaaaac cttagaactc aatgtactgg ttctccagc tctccatcc 600
 tgccgtctcc aggggtgtgcc ccagtgtggg gcaaactgga ccctgagctg ccagtctcca 660
 aggagtaagc ccgctgtcca ataccagtgg gatcggcagc ttccatcctt ccagacttcc 720
 tttgcaccag cattagatgt catccgtggg tctttaagcc tcaccaacct ttcgtcttcc 780
 atggctggag tctatgtctg caaggcccac aatgaggtgg gcaactgcca atgtaatgtg 840
 acgctggaag tgagcacagg tcagttaggg ggcctggagc tgcagtgggt gctggagctg 900
 ttgtgggtac cctggttggg ctgggggtgc tggctgggct ggctctcttg taccaccgcc 960
 ggggcaaggc cctggaggag ccagccaatg atatcaagga ggatgccatt gctccccgga 1020
 ccctgccctg gcccaagagc tcagacacaa tctccaagaa tgggaccctt tctctgtca 1080
 cctccgcacg agccctccgg ccaccccatg gccctccag gccctgggca ttgaccccca 1140
 cgcccagtct ctccagccag gccctgccct caccaagact gcccacgaca gatggggccc 1200
 accctcaacc aatatcccc atccctgggt ggggtttctt ctctggcttg agccgcatgg 1260
 gtgctgtgcc tgtgatgggt cctgcccaga gtcaagctgg ctctctggtg tgatgacccc 1320
 accactcatt ggctaaagga tttgggtctc ctcttcta taagggtcac ctctagcaca 1380
 gaggcctgag tcatgggaaa gagtacact cctgaccctt agtaactctg cccacacctt 1440
 ctttactgtg ggaaaacat ctcatgaaga cctaagtgtc caggagacag aaggagaaga 1500
 ggaagtggat ctggaattgg gaggagcctc caccacccc tgactcctcc ttatgaagcc 1560
 agctgctgaa attagctact caccaagagt gaggggcaga gacttccagt cactgagctt 1620
 cccagggccc cttgatctgt accccacccc tatctaacac cacccttggc tcccactcca 1680
 gctccctgta ttgatataac ctgtcaggct ggcttgggta ggttttactg gggcagagga 1740
 tagggaatct cttattaaaa ctaacatgaa atatgtgttg ttttcatttg caaatttaaa 1800
 taaagataca taatgtttgt atgagataag a 1831

<210> 13
 <211> 909
 <212> DNA
 <213> HOMO SAPIENS

<400> 13
 gaggaggtgg ggcaccaacag acaggcgatt aatgcggctc ttaccaggcc aaccaggact 60
 acagtataca ttgtggacat tcaggacata gattctgcag ctccgggccc acctcactcc 120
 tacctcgatg cctactttgt cttccccaat gggtcagccc tgacccttga tgagctgagt 180
 gtgatgatcc ggaatgatca ggactcgtg acgcagctgc tgcagctggg gctgggtggtg 240
 ctgggctccc agggagagcca ggagtcagac ctgtcgaaac agctcatcag tgtcatcata 300
 ggattgggag tggctttgct gctggtcctt gtgatcatga ccatggcctt cgtgtgtgtg 360
 cggaagagct acaaccggaa gcttcaagct atgaaggctg ccaaggaggc cagggaagaca 420
 gcagcagggg tgatgccctc agcccctgcc atcccaggga ctaacatgta caacactgag 480
 cgagccaacc ccagtctgaa cctccccaac aaagacctgg gcttggagta cctctctccc 540
 tccaatgacc tggactctgt cagcgtcaac tccctggacg acaactctgt ggatgtggac 600
 aagaacagtc agggaaatcaa ggagcacagg ccaccacaca caccaccaga gccagatcca 660
 gagccccctga gcgtggtcct gttaggacgg caggcaggcg caagtggaca gctggagggg 720
 ccactcctaca cctggacctg cctggacacc acggacctgt gacaggggcc cccactcttc 780
 tggacccttt gaagaggccc taccacaccc taactgcacc tgtctccctg gagatgaaaa 840
 tatatgacgc tgcctgcct cctgcttttg gccaatcacg gcagacaggg gttggggaaa 900
 tattttatt 909

<210> 14
 <211> 1453
 <212> DNA

<213> HOMO SAPIENS

<222> 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935

<400> 14

ggagccaagt	ggggccctcg	gcctcttctc	tcgttccagg	cccatgattt	tcctacact	60
tctccctggc	ccaggtctca	gccacaggca	cctctcctgc	ccccgccac	cctcctgacc	120
gcagctccca	ggccctggag	acctccaggc	tttcttgccc	tgggcagccc	cacctcacag	180
ccagagtcaa	tgccttcatg	ggaagggctc	ccagccacac	ccagagtggc	ccaaagctgt	240
tgaagtccag	atcctttgtc	ccatcaggac	cctcctgcct	cctctccagg	cccttggtcg	300
ctccccacc	ctcctcagag	gcccggggaa	gggaagagca	ggtcagtaca	gaggttctgt	360
ctacagggag	gggcccctgg	tctatgcaca	gctggagctc	tgagccttcc	acagcccgtg	420
tgactgctag	agggcagggg	tgcagggctc	agggggggccg	ggctggtcct	ttggggctgg	480
tgttccctacg	tcagtcacca	cctgggggaat	aaactccagc	ctctcctgct	catacagaag	540
gaactggttg	ggtttgcttt	atgggatctt	tgagacaaaa	acagatgctc	ctgtttgctg	600
ggggagggtg	tgagcacgga	gtatttctgt	ccctcgtgaa	gtcacgtcac	acaggggaga	660
ggcgaggtcg	atggaactgg	ccacgcacag	gctctggctc	tggaaggagg	gatgatgagt	720
gggcgttttc	ccggcaggcc	cccggggctc	tcagcctcag	caaccagggg	agaggacaga	780
aatgaaccga	tggttgaggg	attgtcacgg	gaggaacatg	acaccggaag	ggactctagg	840
tgccctcgga	gtgccacaca	tgcccagacc	ttctcacacc	cacacaaata	ggctctgccg	900
tgnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnttggt	cacactcaga	acccaggaca	960
gccacagcca	ccgcttaggg	gaagccactg	cagatgcccc	tggaatgggc	acagcacagc	1020
cagggcgctc	ttccaggcag	gcgaggataa	cttgagagtt	tcctagggca	ccagggacag	1080
agctcagagg	ccccgaggt	gtgtgtagga	ggcggaggcc	cgcagagcac	agagcaggag	1140
aagggcttgg	gccctggagg	agaaaagccat	tctggacacc	aggggacctg	gacggagggt	1200
ccccacagcc	cgtgccccac	gccgcctgga	ggccagaggg	gtcagtggcc	ctgctgtccc	1260
ggctccatct	tggttctagc	cgcacactgt	atgaacacag	tgcccgggt	taacgcacta	1320
acccagcctc	tccctgtgtc	ccacagggag	tagcaagacc	cacccacac	tgcttcacc	1380
atctacacca	gtgacgccgc	tgtgtgtctt	agcatggaaa	taaataaacc	tgaatgcaaa	1440
aaaaaaaaaa	agg					1453

<210> 15
 <211> 443
 <212> DNA
 <213> HOMO SAPIENS

<400> 15

gatatagaca	acttccagag	tcaccagtgt	gcaaattggag	ccacctgcat	tagtcatact	60
aatggctatt	cttgccctctg	ttttggaaat	tttacaggaa	aattttgcag	acagagcaga	120
ttaccctcaa	cagtctgttg	gaatgagaag	acaaatctca	cttgctacaa	tggaggcaac	180
tgacacagagt	tccagactga	attaaaatgt	atgtgccggc	caggttttac	tggagaatgg	240
tgtgaaaagg	acattgatga	gtgtgcctct	gatccgtgtg	tcaatggagg	tctgtgccag	300
gacttactca	acaaattcca	gtgcctctgt	gatgttgcct	ttgctggcga	gcgctgcgag	360
gtggacttgg	cagatgactt	gatctccgac	attttcacca	ctattggctc	agtgactgtc	420
gccttggtac	tgatctctt	gct				443

<210> 16
 <211> 1537
 <212> DNA
 <213> HOMO SAPIENS
 <222> 284, 285, 287

<400> 16

aaaaaaaaaa	acccggtagc	attgtccctt	ccccactgac	aaacttatca	aatccagaag	60
cttttagagtt	tcgtctctaa	ttatttttct	cctgaacaaa	attacccaag	tcaaaacaaa	120
atgtattttt	agaattacgg	cagcatcacga	cctgaatttt	gtgagtttcg	tggttttatc	180
ttaaatacacc	atttccctaa	aaacggtttc	tttctcctta	gaaatgctgg	tggcaacttg	240
atgaaacagc	caaatgcacc	agggcaggtc	actttcccaa	aaannanaag	aaaaaaaaact	300
cattgagata	gctacagttc	tatagggtta	tttaaagcct	cctttttcta	ctcatttttg	360

aaagcaaaat	tacattttac	tattttacat	aaccagtga	aagacgttga	aagcctacag	420
ctcactgttt	ttggtgctct	ggaaatgttg	aggggtgggt	tttaaccagt	gatttttaac	480
gtgcagtga	tttgtagac	ttttaaacac	cagctaaggt	agtcaaactt	gatccccatt	540
aaaaatcaag	gaattagggg	tcgggggagg	gttttaggagt	gatccagaat	gacctcccag	600
aattactgtg	cgtacaactt	tatttttcag	agttttcatt	ggaatggtaa	gagttttatg	660
aaagacagtt	ttaaaactta	ttctgagtta	aatattaata	ctttaaaaaa	ttattgtact	720
agacttattg	cagccttttg	aaagtagcag	agtttcatca	taccacatat	ataacagagc	780
ataaattttc	tataatcagg	caccttttgc	tgcttttgag	taagactggt	ttcctgttta	840
ggtgttaagc	atcgccagac	ataaaaaatct	attctctcct	ctcgattgta	gcatagcctg	900
acagctctag	atacagcatt	tctatgatga	aaaatgagta	tccatcagga	aatctagaag	960
actagccgtg	ttttctcaga	ctccaccttt	gtttgcactc	tggtgcctgt	gaggagcttt	1020
ctggcatgtg	attattttact	tcaaaaactag	agttccaagc	acctacatta	attattttat	1080
attgtgtgca	gaatagtata	tcttttaatg	tcagatatga	tacactgcac	atattgcttt	1140
tgcaactctta	aaatttttgt	actaaataat	agaaaaatatt	tatatctctt	gagtgtgagc	1200
tttgaataga	tggcattatc	actttattgt	ttttttaaca	aaaacttttt	ctcaattatt	1260
ctattgcaat	gttattctga	gcaagtccta	tgccaaatat	cttgataaat	gtttgtatgg	1320
aagattaaat	tttactcttg	tgtggtaaga	ctatttcagt	tactgatttt	atagtgggaa	1380
tttgatattc	cagcacaaag	tccacagtgt	attcagaaat	ccaagttggg	gtcatacatt	1440
tcattttgat	gtgaactttt	ctttgctttc	ctttgttcta	agactccatt	ttgcaataaa	1500
cgttttgaca	gtaaaaaaa	taaaaaagga	aaaaaaa			1537

<210> 17
 <211> 972
 <212> DNA
 <213> HOMO SAPIENS

<400> 17						
acgcaaatc	ggcagcaggg	ttctaaaacc	cagtttggtt	tacgttgtct	ttcacagtag	60
tatatattagc	tcttctctgg	aaagtgtgtg	gttaatatata	ttcttaaaca	tgaaaatgta	120
attaaacaca	ccacgagaga	acaatatctc	aggagactta	atagtgatta	ctttcttcaa	180
tcaggaaatc	gtttcagtc	ctcctttgta	ggaatgcttt	gttttgtgat	gggttttctt	240
aaagaagagc	acacctccgt	ccaatctcct	gagacagcca	cgtctccgct	gacatccac	300
tgatgctct	tcagatagtc	agtgaatggt	tctgataacc	ttcatccagt	atctgaaaca	360
caatgtgaga	gatttatattg	ttttagataa	taacatccca	tttagttgac	taaaatcttc	420
caaactctga	aagctgcaca	ctgctactcc	agagagtgca	ggtcttagct	cttctccttt	480
ctgacttcaa	gatgaatctt	tgggacgatg	tttctgggtg	ttggtccaca	gtgattcact	540
tttgaaggag	aggccacatg	acatgaactg	cctgggtgta	caacctagct	aacatatttg	600
atgctactcc	tggtgtctgt	actgcttatt	caagtagtat	tctaagttat	gttactaaaa	660
aacatggtgg	gtaaagcaca	atcctaccca	tcattgtcct	ccaaaataat	tgtatgacat	720
acacggccca	gccatttgcc	ctccctgcat	ctctgtgetg	ctttgccatt	tccccttcta	780
cccagcctcc	tcaaggggta	ccttggtgga	tatttcagta	cttaaaacca	gactgtaatc	840
ataacctccc	tctgtgtggc	atcaataaat	agccaaactc	aaaaaaaaaa	aaaaaaaaaa	900
aaaaaaaaaa	aaaaaatatc	ggtcgcaagc	ttattccctt	tagtgagggg	taatttttagc	960
ttgcactgcc	ta					972

<210> 18
 <211> 1544
 <212> DNA
 <213> HOMO SAPIENS

<400> 18						
tactttgact	ttggatcatt	tccctgactg	ggctaagtgt	acacatattg	agacttagga	60
agagccacaa	gaccacacac	acagccctta	ccctcctcag	gactaccgaa	ccttctggca	120
caccttgtac	agagtttttg	ggttcacacc	ccaaaatgac	ccaacgatgt	ccacacacca	180
ccaaaaccca	gccaatgggc	cacctcttcc	tccaagccca	gatgcagaga	tggacatggg	240
cagctggagg	gtaggctcag	aaatgaaggg	aaccctcag	tgggctgctg	gacctatctt	300
tcccaagcct	tgccattatc	tctgtgaggg	aggccaggta	gccgagggat	caggatgcag	360
gctgctgtac	ccgctctgcc	tcaagcatcc	cccacacagg	gctctgggtt	tactctgctt	420
cgtcctagat	agttttaaatg	ggaatcagat	cccctgggtg	agagctaaga	caaccaccta	480
ccagtgcacca	tgtcccttcc	agtcacctt	gagcagcctc	agatcatctc	tgtcactctg	540
gaaggacac	ccagccagg	gacggaatgc	ctggctctga	gcaacctccc	actgctggag	600
tgcgagtggg	aatcagagcc	tctgaagcc	tctgggaact	cctcctgtgg	ccaccaccaa	660
aggatgagga	atctgagttg	ccaacttcag	gacgacacct	ggcttgccac	ccacagtgca	720

ccacaggcca	acctacgccc	ttcatcactt	ggttctgttt	taatcgactg	gccccctgtc	780
ccacctctcc	agtgagcctc	cttcaactcc	ttggtccctt	gttgtctggg	tcaacatttg	840
ccgagacgcc	ttggctggca	ccctctgggg	tccccctttt	ctcccaggca	ggtcatcttt	900
tctgggagat	gcttcccttg	ccatcccca	atagctagga	tcacactcca	agtatgggca	960
gtgatggcgc	tctggggggc	acagtgggct	atctagggtc	tccctcacct	gaggcccaga	1020
gtggacacag	ctgttaattt	ccactggcta	tgccacttca	gagtctttca	tgccagcggt	1080
tgagctcctc	tgggtaaaat	cttccctttg	ttgactggcc	ttcacagcca	tggctgggtga	1140
caacagagga	tcgttgagat	tgagcagcgc	ttggtgatct	ctcagcaaac	aaccctgcc	1200
cgtgggcca	tctacttgaa	gttactcgga	caaagacccc	aaagtggggc	aacaactcca	1260
gagaggctgt	gggaatcttc	agaagcccc	ctgtaagaga	cagacatgag	agacaagcat	1320
cttctttccc	ccgcaagtcc	attttatatt	cttcttgtgc	tgctctggaa	gagaggcagt	1380
agcaaagaga	tgagctcctg	gatggcattt	tccagggcag	gagaaagtat	gagagcctca	1440
ggaaccccca	tcaaggaccg	agtatgtgtc	tgggttccttg	ggtgggacga	ttcctgacca	1500
actgtccag	ctcttgctct	cattaaatgc	tctgtctccc	gcgg		1544

<210> 19

<211> 1109

<212> DNA

<213> HOMO SAPIENS

<222> 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 884, 885

<400> 19

cggacgcgtg	ggggccagcc	tggaggccca	gacgtggcgc	agcgactcgg	aggttcgcct	60
ccagcttgcg	catcatctgc	ggccgggtcc	cgatgagcct	cctgttgccct	ccgctggcgc	120
tgtgtctgct	tctcgcggcg	cttgtggccc	cagccacagc	cgccactgcc	taccggccgg	180
actggaaccg	cttgagcggc	ctaaccgcgc	ccgggtaga	gacctgcggg	ggatgacagc	240
tgaaccgcct	aaaggaggtg	aaggctttcg	tcacgcagga	cattccattc	tatcacaacc	300
tgggtgatgaa	acacctccct	ggggccgacc	ctgagctcgt	gctgctgggc	cgccgctacg	360
aggaactaga	gcgcatccca	ctcagtga	tgaccgcgca	agagatcaat	gcgctagtgc	420
aggagctcgg	cttctaccgc	aaggcggcgc	ccgacgcgca	ggtgcccccc	gagtacgtgt	480
ggggcggcgc	gaagcccca	gaggaaaact	cggaccacgc	tgacctgtag	gtccgggggc	540
gcggcggagc	tgggacctac	ctgcctgagt	ctgggagaca	gaatgaagcg	ctcagcatcc	600
cggaataact	tctcttctgc	agagccgatg	cccgctcccc	ggccagcagg	gatgggggtg	660
gggaggttct	cccaacccca	cttctctcct	tccccagctc	cactaaattc	cctcctgcct	720
taaaaaaac	aagaaaaacc	aaacaaacaa	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	780
nnnnnnnnnn	nnnnnnnnnn	nnnnnnntct	ctatagtgtc	acctaaattc	aattcactgg	840
ccgtcgtttt	acaacgtcgt	gactggcgac	ggacaaagtt	atcnntttta	tcgccttgca	900
gcacataccc	ctttggccag	ctggggtaat	aggggaagcg	ggccggaccc	gatcggcctt	960
cccaaacagt	tggggaagct	tgaaatgcgc	gacattgggc	cgacggcctt	ctatacggga	1020
ggatctctaa	acgcggccgg	ggtgttgggt	gggttaaggc	ggagtgtgac	cccgcataat	1080
aacttttgca	caggggcccct	atagggggcc				1109

<210> 20

<211> 1740

<212> DNA

<213> HOMO SAPIENS

<400> 20

aagagaagtt	accccgatga	cttgggtttgg	aaggggttaa	ggcaccagtg	catcctcttc	60
taaagtgatt	tatgatgatg	tgtggagtgt	aaaaacttta	ccccacccca	aagaacagcc	120
ctctcactcc	tactgagtc	cactctgaac	gtgctaaaat	gggaaggagg	cggtgttttg	180
ctgatctgtt	aaattcttag	tgaagtcttc	ttgatttcca	gtggctgctg	ttgtttgagt	240
ttggtttggg	gcaaaactga	ggtagtccca	acatttctgg	gactgaatcc	aggcaagaga	300
aagaagaaaa	agaagaagaa	aaagaggagg	aaaaaggtag	ggagaaataa	agggaggaga	360
gaagcacagt	gaaagaaaaa	aaaagtcctt	tttgcgacat	cacattcctg	tgttttccct	420
cagcctggaa	aacatattaa	tcccagtgtc	tttacgcccc	gaaacaaaga	gactaagcca	480
gactatgggg	gaaagggaga	taagaaggat	cctggaaact	taaagaggga	aagagtggga	540
ttcagaaatc	gccaggactg	gactttaagg	gacgtcctgt	gtcagcacia	gggactggca	600
cacacagaca	cacgagaccg	aggagaaact	gcagacaaat	ggagatacaa	agacttagaa	660
ggacagctcc	tttcacctca	tcctacttgt	ccagaaggta	aaaagacaca	gccagaaaga	720

aaagggcatcg	gctcagctct	cagatcagga	caggctgtgg	atctgtggcg	gtactctgaa	780
agctggagct	gcagcacacc	ccttttgtat	tgctcaccct	cggtaaagag	agagagggct	840
gggaggaaaa	gtagtctatc	taggaaactg	tcctgggaac	caaacttctg	atttcttttg	900
caaccctctg	cattccatct	ctatgagcca	ccattggatt	acacaatgac	atggagaatg	960
ggaccccggt	tcactatgct	gttggccatg	tggctagtgt	gtggatcaga	acccaccccc	1020
catgccacta	ttagaggcag	ccacggagga	cggaaagtgc	ctttggtttc	tccggacagc	1080
agtagggcag	ctcggtttct	gaggcacact	gggaggtctc	gcggaattga	gagatccact	1140
ctggaggaaac	caaaccttca	gcctctccag	agaaggagga	gtgtgcccgt	gttgagacta	1200
gctcgcccaa	cagagccgcc	agcccgctcg	gacatcaatg	gggccgccgt	gagacctgag	1260
caaagaccag	cagccagggg	ctctccgcgt	gagatgatca	gagatgaggg	gtcctcagct	1320
cgggtcaagaa	tgttgcgttt	cccttacggg	gtccagctct	cccaacatcc	ttgccagctt	1380
tgcagggaag	aacagagtat	gggtcatctc	agccctcat	gcctcggaag	gctactaccg	1440
cctcatgatg	agcctgctga	aggacgatgt	gtactgtgag	ctggcggaga	ggcacatcca	1500
acagattgtg	ctcttcacc	aggcaggtga	ggaaggaggc	aagggtgagaa	ggatcaccag	1560
cgagggccag	atcctggagc	agccctggg	accctagcct	catccctaag	ctgatgagct	1620
tcctgaagct	ggagaagggc	aagtttggca	tgggtgctgt	gaagaagacg	ctgcagggtg	1680
aggagcgcta	tccatatccc	gttaggctgg	aagccatgta	cgagggtcatc	gaccaaggcc	1740

<210> 21

<211> 4467

<212> DNA

<213> HOMO SAPIENS

<222> 971, 978, 1295, 1296, 1297, 1298, 1299, 1300, 1301, 1302, 1303, 1304, 1305, 1306, 1307, 1308, 1309, 1310, 1311, 1312, 1313, 1314, 1315, 1316, 1317, 1318, 1319, 1320, 1321, 1322, 1323, 1324, 1325, 1326, 1327, 1328, 1329, 1330, 1331, 1332, 1333, 1334, 1335, 1336, 1337, 1338, 1339, 1340, 1341, 1342, 1343, 1344, 1345, 1346, 1347, 1348, 1349, 1350, 1351, 1352, 1353, 1354, 1355, 1356, 1357, 1358, 1359, 1360, 1361, 1362, 1363, 1364, 1365, 1366, 1367, 1368, 1369, 1370, 1371, 1372, 1373, 1374, 1375, 1376, 1377, 1378, 1379, 1380, 1381, 1382, 1383, 1384, 1385, 1386, 1387, 1388, 1389, 1390, 1391, 1392, 1393, 1394, 1395, 1396, 1397, 1398, 1399, 1400, 1401, 1402, 1403, 1404, 1405, 1406, 1407, 1408, 1409, 1410, 1411, 1412, 1413, 1414, 1415, 1416, 1417, 1418, 1419, 1420, 1421, 1422, 1423, 1424, 1425, 1426, 1427, 1428, 1429, 1430, 1431, 1432, 1433, 1434, 1435, 1436, 1437, 1438, 1439, 1440, 1441, 1442, 1443, 1444, 1445, 1446, 1447, 1448, 1449, 1450, 1451, 1452, 1453, 1454, 1455, 1456, 1457, 1458, 1459, 1460, 1461, 1462, 1463, 1464, 1465, 1466, 1467, 1468, 1469, 1470, 1471, 1472, 1473, 1474, 1475, 1476, 1477, 1478, 1479, 1480, 1481, 1482, 1483, 1484, 1485, 1486, 1487, 1488, 1489, 1490, 1491, 1492, 1493, 1494, 1495, 1496, 1497, 1498, 1499, 1500, 1501, 1502, 1503, 1504, 1505, 1506, 1507, 1508, 1509, 1510, 1511, 1512, 1513, 1514, 1515, 1516, 1517, 1518, 1519, 1520, 1521, 1522

<400> 21

gcgtgcgct	caccctgcgc	gtgccccgc	ggcgctggg	cgtcttctg	gactacgagg	60
ccggagagct	gtccttcttc	aacgtgtccg	acggctcca	catcttcacc	ttccacgaca	120
ccttctcg	cgcgctctgt	gcgtacttca	ggcccagggc	ccacgacggc	ggcgaacatc	180
cggatcccc	gaccatctgc	ccgctgccgg	ttagagggac	gcgcgtcccc	gaagagaacg	240
acagtgcac	ctggctacag	ccctatgagc	ccgcggatcc	cgccctggac	tgggtgtgag	300
gcgcctcgt	ggcgcgggga	ctggccccgc	ggggggcccc	tggatcccag	gccagcgctt	360
tgctctcctg	ctccgtctga	agggagcagg	tgcaccagcc	aaaatgtcag	cgaggggggac	420
aaagagaggg	acctttgcct	acgtagatgt	gtatgtgtag	tgcgattttc	ttcaaggaaa	480
ggagacaagt	ccaaagctcg	tttgtggatt	gtgggactga	gcaaaggagt	acaaatatat	540
ccacgtcgct	cagagctggg	gtgctcacgg	tgggtgggtg	gaaagaagcc	agcatgggaag	600
aaagaaggga	gaaaactttg	gtgactgcct	tagagggatc	agttaatttg	tatagtttta	660
tattttttgt	atatgtttgc	tagctctaaa	aagggtcgaga	tgcaataaca	cttcgtaagc	720
aacgagttca	cctaagtaag	gctcagatcc	tagtttttaa	aaccatttcc	cattaaaatg	780
aagttggagg	aacagctgct	tctggagccg	gggcaaaaaa	tttcaagggtg	agcctggagc	840
attgtgtgtg	gtgaagtaaa	ataaaggctc	aaaacgtgac	ggcaacccgg	caaaagggtg	900
gggagccagg	ccgaagggcc	tcactgacca	attgtgggac	aatttgaaca	tcaggatgaa	960
taatgcacag	ngaggttnta	acacactgaa	taaaaacata	atccatgagt	tcatgtctgat	1020
actcaaattt	cttttttaaaa	aggagaaaca	ggaagggttc	ttttggaggt	gaaatctaata	1080
tattggtgag	agtcttggag	aacaggctgt	ttccagtcctc	aaagcagtaa	ccttatacac	1140

tacttataag	tttgaaaggg	gaaagggtac	ctttacaatg	gagacatcta	ccagatgcat	1200
ccaagtgatt	aaatttaaca	tcatcaatga	tgggaccaag	gacattatta	gtttgacaac	1260
tggggaaaga	agtgttcttc	acccctacc	ccannnnnn	nnnnnnnnnn	nnnnnnnnnn	1320
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	1380
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	1440
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	1500
nnnnnnnnnn	nnnnnnnnnn	nnttcatttt	cagagtga	catttgtact	gtggctatgt	1560
aggagaacat	tcttgttctt	agcaaacata	ctgaagtttt	tagatattaa	ttaccacagt	1620
gtctgccact	gaatttccag	tgactaagtg	gaaaaatata	aaacatatga	atataaagaa	1680
agaaagagac	aagtcaaagt	tagtaaaatg	acaacacttg	gtgactctag	gtgactggtc	1740
gacagatgtt	cattgtacta	tcaatgtggc	tttgtgtggg	gtttgaaatt	ttgcaaacta	1800
agagttgggt	ggcggggaga	aggatacacc	aaaaaactaa	gtgattatct	ttggatggga	1860
aaatgttttg	taattgcatt	cttaaaatgt	cttcttttga	ttttttaatg	ttcaataatg	1920
tatatgtatc	agttctgtaa	taaaggggaa	aacacttttt	ttaaatactc	ataaaaaacc	1980
atccgtagga	tcgagaagat	caggcagaag	ggctttgtcc	agaaatgtaa	ggcctctggt	2040
gtagagggcc	aggtggtggc	ggaggggaat	gacggtggag	ggggagcagg	aaggccaagc	2100
ctgggcagcg	agaagaagaa	agaggaccca	aggagagcac	aagtccacc	aaccagagag	2160
agtcgggtga	aggtcctgag	aaaactggcc	gccactgcac	cagctttgcc	ccaacctccc	2220
tcaaccccca	gagccaccac	ccttctctct	gccccaggcc	acaacagtga	ctcgggtccac	2280
gtcccgggcg	gtaacagttg	ctgcaagacc	tatgaccacc	actgcctttc	ccaccacggc	2340
agaggccctg	gacccctca	ccctcccaca	ggccccctac	aaccactgag	gtgatcactg	2400
ccaggagacc	ctcagtttca	gagaatcttt	accctccatc	ccggaaggat	cagcacaggg	2460
agaggccaca	gacaaccagg	aggcccagca	aggccaccag	cttgagagag	ttcacaaatg	2520
cccctcccac	caccatctca	gaaccagca	caagggtgct	tggcccaggc	cgtttccggg	2580
acaaccgcat	ggacaggcgg	gaacatggcc	accgagaccc	aaatgtgggt	ccaggctcctc	2640
ccaagccagc	aaaggagaaa	cctcccaaaa	agaaggccca	ggacaaaatt	cttagtaatg	2700
agtatgagga	gaagtatgac	ctcagccggc	ctactgcctc	tcagctggag	gacgagctgc	2760
aggtggggaa	tgttccccctt	aaaaaagcaa	aggagtctaa	aaagcatgaa	aagcttgaga	2820
aaccagagaa	ggagaagaaa	aaaaagatga	agaatgagaa	cgcagacaag	ttacttaaga	2880
gtgaaaagca	aatgaagaag	tctgagaaaa	agagcaagca	agagaaagag	aagagcaaga	2940
agaaaaaagg	aggtaaaaaca	gaacaggatg	gctatcagaa	accaccaac	aaacacttca	3000
cgcagagtcc	caagaagtca	gtggccgacc	tgctgggggtc	ctttgaaggc	aaacgaagac	3060
tccttctgat	cactgctccc	aaggctgaga	acaatatgta	tgtgcaacaa	cgtgatgaat	3120
atctggaag	tttctgcaag	atggctacca	ggaaaatctc	tgtgatcacc	atcttcggcc	3180
ctgtcaacaa	cagcaccatg	aaaatcgacc	actttcagct	agataatgag	aagcccatgc	3240
gagtgggtgga	tgatgaagac	ttggtagacc	aagcgtctca	tcagcgagct	gaggaaagag	3300
tacggaatga	cctacaatga	cttcttcatg	gtgctaacag	atgtggatct	gagagtcaag	3360
caatactatg	aggtaccaat	aacaatgaag	tctgtgtttg	atctgatcga	tactttccag	3420
tcccgaatca	aagatatgga	gaagcagaag	aaggagggca	ttgtttgcaa	agaggacaaa	3480
aagcagtcct	tggagaactt	cctatccagg	ttccggtgga	ggaggagggt	gctgggtgatc	3540
tctgtccta	acgatgaaga	ctgggcctat	tcacagcagc	tctctgccct	cagtggctcag	3600
gcgtgcaatt	ttggtctgcg	ccacataacc	attctgaagc	ttttaggcgt	tggagaggaa	3660
gttgggggag	tgttagaact	gttcccaatt	aatgggagct	ctgttgttga	gcgagaagac	3720
gtaccagccc	atttggtgaa	agacattcgt	aactattttc	aagtgagccc	ggagtacttc	3780
tccatgcttc	tagtcggaaa	agacggaaat	tgc aaatctc	ggatatcctc	cccaatgtgg	3840
tccatgggtga	ttgtgtacga	tttaattgat	tcgatgcaac	ttcggagaca	ggaaatggcg	3900
attcagcagt	cactggggat	gcgctgcccc	gaagatgagt	atgcaggcta	tggttaccat	3960
agttaccacc	aaggatacca	ggatgggttac	caggatgact	accgtcatca	tgagagttat	4020
caccatggat	acccttactg	agcagaaata	tgtaacctta	gactcagcca	gtttcctctg	4080
cagctgctaa	aactacatgt	ggccagctcc	attcttccac	actgcgtact	acatttccctg	4140
cctttttctt	tcagtgtttt	tctaagacta	aataaaatgc	aaactttcac	ctattctatga	4200
gttattattg	aaacctcaaa	tcataaagac	attttaaaga	attgtttttc	taactggagg	4260
ggctctagt	ctaaataata	gtactgaaaa	ttgatattat	tttctttttc	ttatatgaag	4320
gaccttattt	ggcatataaa	attttataaa	atatgtattt	aaagcttttt	cttatttttt	4380
gtattaattg	gtaagtga	actctgttaa	agatcacacc	acaatgtttt	caagaaacat	4440
ctgaaaagat	aaaacaaaga	acaaata				4467

<210> 22

<211> 2965

<212> DNA

<213> HOMO SAPIENS

<222> 1473, 1474, 1475, 1476, 1477, 1478, 1479, 1480, 1481, 1482, 1483, 1484, 1485, 1486, 1487, 1488, 1489, 1490, 1491, 1492, 1493, 1494, 1495,

1496, 1497, 1498, 1499, 1500, 1501, 1502, 1503, 1504, 1505, 1506, 1507,
 1508, 1509, 1510, 1511, 1512, 1513, 1514, 1515, 1516, 1517, 1518, 1519,
 1520, 1521, 1522, 1523, 1524, 1525, 1526, 1527, 1528, 1529, 1530, 1531,
 1532, 1533, 1534, 1535, 1536, 1537, 1538, 1539, 1540, 1541, 1542, 1543,
 1544, 1545, 1546, 1547, 1548, 1549, 1550, 1551, 1552, 1553, 1554, 1555,
 1556, 1557, 1558, 1559, 1560, 1561, 1562, 1563, 1564, 1565, 1566, 1567,
 1568, 1569, 1570, 1571, 1572, 1573, 1574, 1575, 1576, 1577, 1578, 1579,
 1580, 1581, 1582, 1583, 1584, 1585, 1586, 1587, 1588, 1589, 1590, 1591,
 1592, 1593, 1594, 1595, 1596, 1597, 1598, 1599, 1600, 1601, 1602, 1603,
 1604, 1605, 1606, 1607, 1608, 1609, 1610, 1611, 1612, 1613, 1614, 1615,
 1616, 1617, 1618, 1619, 1620, 1621, 1622, 1623, 1624, 1625, 1626, 1627,
 1628, 1629, 1630, 1631, 1632, 1633, 1634, 1635, 1636, 1637, 1638, 1639,
 1640, 1641, 1642, 1643, 1644, 1645, 1646, 1647, 1648, 1649, 1650, 1651,
 1652, 1653, 1654, 1655, 1656, 1657, 1658, 1659, 1660, 1661, 1662, 1663,
 1664, 1665, 1666, 1667, 1668, 1669, 1670, 1671, 1672, 1673, 1674, 1675,
 1676, 1677, 1678, 1679, 1680, 1681, 1682, 1683, 1684, 1685, 1686, 1687,
 1688, 1689, 1690, 1691, 1692, 1693, 1694, 1695, 1696, 1697, 1698, 1699,
 1700, 1701, 1702, 1703, 1704, 1705, 1706, 1707, 1708, 1709, 1710, 1711,
 1712, 1713, 1714, 1715, 1716, 1717, 1718, 1719, 1720, 1721, 1722, 1723,
 1724, 1725, 1726, 1727, 1728, 1729, 1730, 1731, 1732, 1733, 1734, 1735,
 1736, 1737, 1738, 1739, 1740, 1741, 1742, 1743, 1744, 1745, 1746, 1747,
 1748, 1749, 1750, 1751, 1752, 1753, 1754, 1755, 1756, 1757, 1758, 1759,
 1760, 1761, 1762, 1763, 1764, 1765, 1766, 1767, 1768, 1769, 1770, 1771,
 1772, 1773, 1774, 1775, 1776, 1777, 2948, 2951, 2961

<400> 22

aaacaaagt	caatttagct	ggattttctga	actatgggtt	tgaatgttta	aagaagaatg	60
atgggtacag	ttaggaaagt	ttttttctta	caccctgac	ttgagggaaa	cattgcttgt	120
ctttgagaaa	ttgactgaca	tactggaaga	gaacaccatt	ttatctcagg	ttagtgaaga	180
atcagtgcag	gtccctgact	cttattttcc	cagaggccat	ggagctgaga	ttgagactag	240
ccttgtggt	ttcacactaa	agagtttcc	tggtatgggc	aacatgcatg	acctaattgtc	300
ttgcaaaatc	caatagaagt	attgcagctt	ccttctctgg	ctcaagggtc	gagttaagt	360
aaaggaaaaa	cagcacaatg	gtgaccactg	ataaaggctt	tattaggtat	atctgaggaa	420
gtgggtcaca	tgaatgttaa	aaagggaatg	agggttttgt	tggttttttg	aagtaaaagg	480
aaacataaat	attaccatga	tgaattctag	tgaatgacc	ccttgacttt	gcttttctta	540
atacagatat	ttactgagag	gaactatttt	tataacacaa	gaaaaattta	caattgatta	600
aaagtatcca	tgtcttggt	acatacgtat	ctatagagct	ggcatgtaat	tcttctctca	660
taaagaatag	gtataggaaa	gactgaataa	aaatggaggg	atatccctct	ggatttcaat	720
tgcattgtgc	aataagcaaa	gaagggttga	taaaagttct	tgatcaaaaa	gttcaaaagg	780
accagaattt	tagacagcaa	gctaaataaa	tattgtaaaa	ttgcactata	ttagggttaag	840
tattatttag	gtattataat	atgctttgta	aattttatat	tccaaatatt	gctcaatatt	900
tttcatctat	taaatttaatt	tctagtataa	ataagtagct	tctatatctg	tcttagtcta	960
ttataattgt	aaggagtaaa	attaaatgaa	tagtctgcag	gtataaattt	gaacaatgca	1020
tagatgatcg	aaaattacgg	aaaatcatag	ggcagagagg	tgtgaagatt	catcattatg	1080
tgaattttgg	atcttttctca	aatccttgct	gaaatttagg	atggttctca	ctgtttttct	1140
gtgctgatag	taccctttcc	aagggtgacct	tcagggggat	taaccttctc	agctcaagca	1200
aggagctaaa	aggagcctta	tgcatgatct	tcccacatat	caaaataact	aaaaggcact	1260
gagtttgga	tttttctgcc	tgctctgcta	agaccttttt	ttttttttac	tttcattata	1320
acataattata	ctgcacatta	tacaaaaatg	attaaaatat	attaaaacaa	catcaacaat	1380
ccaggatatt	tttctataaa	acttttttaa	aataattgta	tctatatatt	caattttaca	1440
tcctttttca	aaggctttgt	ttttctaaag	gcnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	1500
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	1560
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	1620
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	1680
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	1740
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	1800
ttcttaattg	ctgcataata	tcacatcaaa	taggcatttt	tcaaacctct	ttccttatta	1860
aacatgtaga	ctatatccat	tttttactaa	aataaataac	atttcagata	atatctttgc	1920
actgataatg	ttgccaagcc	atttctaaag	tgaccttatt	aatttaatta	ccattggatg	1980
agggtgttgc	tttcatcgca	ccattgtaga	ttgtcttttt	tatttcaatt	tgcgtttatt	2040
tataactggt	tgcaaaagga	cacagaacac	acgtctcttc	aacttatctt	tgataaaacc	2100
aagcaaggat	gcaaaaagtt	ggacgacatt	gagtagagtc	atgggtatac	gtgctgacct	2160
tacagtatca	gtggaaaaga	taaggaaaat	gtcactactc	acctatgtta	tgcaaaacag	2220

ttaggtgtgc	tggggctgga	tactgctctt	ttacttgagc	attggttgat	taaagtttag	2280
gtaccatcca	gggctgggtct	agagaagtct	ttggagttaa	ccatgctctt	tttggttaaag	2340
aagagagtaa	tgtgtttatc	ctggctcata	gtccgtcacc	gaaaatagaa	aatgccatcc	2400
ataggtaaaa	tgctgacct	tagaaaaaaa	tgaactctac	ttttatagcc	tagtaaaaaat	2460
gctctacctg	agtagttaaa	agcaattcat	gaagcctgaa	gctaaagagc	actctgatgg	2520
ttttggcata	atagctgcat	ttccagacct	gacctttggc	cccaaccaca	agtgtctcaa	2580
gccccaccag	ctgaccaaaag	aaagcccaag	ttctccttct	gtccttccca	caacctccct	2640
gtccccaaaa	ctatgaaatt	aatttgacca	tattaacaca	gctgactcct	ccagtttact	2700
taaggtagaa	agaatgagtt	tacaacagat	gaaaataagt	gctttgggcg	aactgtattc	2760
cttttaacag	atccaaacta	ttttacattt	aaaaaaaaag	ttaaactaaa	cttctttact	2820
gctgatatgt	ttcctgtatt	ctagaaaaat	ttttacactt	tcacattatt	tttgtacact	2880
ttccccatgt	taagggatga	tggcttttat	aaatgtgtat	tcattaaatg	ttacttttaa	2940
aataaaaana	naaaaaaaaa	naaaa				2965

<210> 23

<211> 1734

<212> DNA

<213> HOMO SAPIENS

<222> 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899

<400> 23

cgccctccgga	aactgcccc	cgggctgctg	gccaaacttca	ccctcctgcg	cacccttgac	60
cttggggaga	accagttgga	gaccttgcca	cctgacctcc	tgagggggcc	gctgcaatta	120
gaacggctac	atctagaagg	caacaaattg	caagtactgg	gaaaagatct	cctcttgccg	180
cagccggacc	tgcgtacct	cttctgaac	ggcaacaagc	tgccagggt	ggcagccggt	240
gccttccagg	gcctgcgga	gctggacatg	ctggacctct	ccaataactc	actggccagc	300
gtgcccagg	ggctctgggc	atccctaggg	cagccaaact	gggacatgcg	ggatggcttc	360
gacatctccg	gcaaccctcg	gatctgtgac	cagaacctga	gcgacctcta	tcgttggctt	420
caggcccaaa	aagacaagat	gttttccag	aatgacacgc	gctgtgctgg	gcctgaagcc	480
gtgaagggcc	agacgtcct	gggcagtgcc	caagtcaccg	tgagaccagg	ggcttgggtt	540
gagggtgggg	ggtctggtag	aacactgcaa	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	600
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	660
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	720
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	780
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	840
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnna	900
taatcctgct	tttacagggt	aaactcgagg	ctgtccatag	cggtctgggac	cccgtttcat	960
ccatccatgc	ttcctagaac	acacgatggg	ctttccttac	ccatgcccaa	ggtgtgccct	1020
ccgtctggaa	tgccgttccc	tgtttcccag	atctcttgaa	ctctgggttc	tcccagcccc	1080
ttgtccttcc	ttccagctga	gccttgccca	cactggggct	gcctttctct	gactctgtct	1140
tcccaagtc	agggggctct	ctgagtgcag	ggtctgatgc	tgagtccac	ttagcttggg	1200

gtcagaacca	aggggttttaa	taaataaacc	ttgaaaactg	gatcggatga	attggctttc	1260
attgtgttcc	tagcatcttc	tcaaatcaac	ttcccaggac	tccagggatga	aggaggaaaa	1320
gagggcatggc	ccaggccctg	gggtgtggga	tatgggtctcc	ctaggggatg	acagttggga	1380
tcaatggcct	gtgacttctc	ctctcccttc	ccccatcctg	ggacctaaact	ggaaaataaaa	1440
ccttgactgt	tgccccgggtg	tcattttacc	agtggatttc	tgccaggggt	tgtgtcctag	1500
gagaaggttt	aagttaaacc	agattgcccc	ggctctccaaa	cgatttgtca	tgctgacctg	1560
agatcatcga	agggggcacc	tgcccccggtg	caagggtgca	ggggcaggat	ggggctgaag	1620
ggatgagcag	gggtcccggtg	ccacctgctg	atacagcatt	ggccatgtgg	gggctgcaat	1680
cggattttgga	agaccctggg	gcttggggggc	atgtccattt	tcccagtcctc	taaa	1734

<210> 24

<211> 4005

<212> DNA

<213> HOMO SAPIENS

<400> 24

ggacaccgtc	tgcaagtggag	tcaactggtgc	cgtaaatgtg	gccaagggggg	ccgtccagac	60
gggctgtaga	cacggccaag	accgtgctga	ccggcaccaa	ggacacagtc	actactgggc	120
tcatgggggc	agtgaatgtc	gccc aaaggga	ccgtccagac	cagtgtggac	accaccaaga	180
ctgtcctaac	tggtaccaag	gacaccgtct	gcagtgggggt	gaccggtgct	gcgaatgtgg	240
ccaagggggc	cgctccagggg	ggcctgggaca	ctacaaagt	tgctcctgact	ggcactaaag	300
acaccgtatc	caactgggctc	acaggggctg	tgaacttggc	caaagggact	gtccagaccg	360
gcgtggacac	cagcaagact	gtcctgaccg	gtaccaagga	caccgtctgc	agtggagtca	420
ctggtgcccgt	aaatgtggcc	aaaggcaccg	tccagacagg	tgtggacaca	gccaagacgg	480
tgctgagtgg	cgctaaggat	gcagtgacta	ctggagtac	gggggacagt	aatgtggcca	540
aagggaaccgt	gcagaccggc	gtggacgcct	ccaaggctgt	gcttatgggt	accaaggaca	600
ctgtcttcag	tggggttacc	ggtgccatga	gcatggccaa	aggggcccgtc	cagggggggcc	660
tggacaccac	caagacagt	ctgaccggaa	ccaaagacgc	agtgtccgct	gggctcatgg	720
ggtcagggaa	cgtggcgaca	ggggccaccc	acactggcct	cagcaccttc	cagaactggt	780
tacctagtac	ccccgccacc	tcctgggggtg	gactcaccag	ttccaggacc	acagacaatg	840
gtggggagca	gactgccctg	agcccccaag	aggccccgtt	ctctggcacc	tccacgcccc	900
cggatgtgct	cagtgtaggc	ccggagcctg	cctgggaagc	cgcagccact	accaagggcc	960
ttgcgactga	cgtggcgacg	ttcacccaag	gggcccgcctc	aggcagggag	gacacggggc	1020
ttttggccac	cacacacggc	cccgaagaag	ccccagcctt	ggcaatgctg	cagaatgagt	1080
tggaggggct	gggggacatc	ttccaccca	tgaatgcgga	ggagcaagct	cagctggctg	1140
cctcccagcc	cggggccaaag	gtgctgtcgg	cggaacaggg	gagctacttc	gttcgttttag	1200
gtgacctggg	tcccagcttc	cgccagcggg	catttgaaca	cgcgggtgagc	cacctgcagc	1260
acggccagtt	ccaagccagg	gacactctgg	cccagctcca	ggactgcttc	aggctgattg	1320
aaaaggccca	gcaggctcca	gaagggcagc	cacgtctgga	ccagggtcca	ggtgcccagt	1380
cggaggacgc	tgtgtccag	gaggagcggg	atgcgggggt	tctgtccagg	gtctgcggcc	1440
ttctccggca	gctgcacacg	gcctacagt	gcctgggtctc	cagcctccag	ggcctgcccc	1500
ccgagctcca	gcagccagt	gggcccggcg	ggcacagcct	ctgtgagctc	tatggcatcg	1560
tggcctcagc	tggctctgta	gaggagctgc	ccgcagagcg	gctggtgcag	agccgcgagg	1620
gtgtgcacca	ggcttggcag	gggttagagc	agctgctgga	gggcttacag	cacaatcccc	1680
cgctcagctg	tgtgttaggg	cccttcgcct	tgcccgtgtg	cgggcagtag	ctgtaggagc	1740
ctgcaggccc	ggcgcgggggt	cgccctgctc	tgtccaggga	ggagctgcct	cagaactttc	1800
tccccgcccc	caaacctgga	tcggttcctt	aaagccctag	acctttgggg	ctgcagctgg	1860
ctgagcgccg	aggggctgcg	gaggcagtg	ccttcttaac	tgagccaccc	cacgcccctgc	1920
tcggggcctg	cctgcattctc	ccacctctc	cccagcgtg	cctgcccctc	tcggagcctg	1980
gggtcactca	aaaaaccagc	caagagcctt	cccttgaagt	ccccaaagcaa	gcactgcaat	2040
taggaaagag	gtgcccagcc	tggaagggca	tctgttttgc	ccgctagcaa	ccgctagcaa	2100
cccttttata	tctagcaggg	ctcttccagt	cctgcagcac	gggccccccag	ctatcagcgg	2160
tgcaggcagt	gctgtggcat	cccaggctcc	gggcagctcc	gttctcatgc	tgaagtgagg	2220
tctccggcct	tagcacacac	accttgaggg	tcttaagaac	cacattccct	catagtagaa	2280
agtactagaa	aaagcgacac	tgccatcatc	atcccaaggc	aggctgtctac	tgcctttgct	2340
gacccccggg	ttggccctcac	ggtggggact	aagctggccag	gagccacagc	agccacagct	2400
ggggcttttg	accagcctgg	cttgagactg	agcagtttgc	aggggggtggg	gggtgcaaaa	2460
aacaagcaaa	caggctgtctg	ctgcctccag	ctgcccacca	caggcctgcc	ccaggcacct	2520
ggggctctga	ggccccctggg	gaggctgggg	ccagcagctg	cccctggaga	acacagacaa	2580
aggacttccc	cgcaggggaa	tgtgccctat	ggaggggatca	gacaggggctg	ggaacagcca	2640
cagaggctgc	gtgcctatgg	cacagccctt	cctccggcgc	acactccccc	tgggtcctca	2700
ggcccaccca	agcgcggggc	tgccagaggaa	gggggctggg	ggaggctgca	ggcatcagag	2760
acactgggtg	tggcgggacc	ggccgcgggg	cccgtgtctc	tcaggctagc	ccaggctcgtg	2820

gaggctggca	ggctcaggtc	gggtgtgaga	cgtgccgtgg	ctgcgctcag	tccagcgggg	2880
aggagccgtt	cagcccggcc	tccccaggaa	gccatatccc	cactcaccgc	gtaagagaac	2940
cttgtcgtcc	cctttccatg	ctctcctagg	acacgagccc	aggaacccca	gaccagggg	3000
gaggaagggg	ggagggggcc	caggggtcac	catgtgcacc	aggggcccgtg	aggggcccggg	3060
gcattcagct	cagctctgaa	ccgggggaagc	tggcacggca	aggactgcct	caggtgacgg	3120
gccgtgagag	gggacgggtc	aggagccttc	ccaagccttc	tcctcagccc	gacacccatg	3180
gccatcggag	gctaggatgc	cagacacagc	catttgcaga	aatcaggcac	agtgactgca	3240
gctcacgtcc	agccaaccaa	gcattggggcc	gcagctcagg	aagtccttc	ccgccacacc	3300
acagccta	tcttactggg	acggaggcaa	ctcgggtacg	ctgggcagga	cgacaaacac	3360
gagacgccac	tgtggaatga	gcaacttcgg	agcacggggg	gacttgcttg	ggaccgtgcc	3420
cacgtgacag	ccccttatgc	agaggaggaa	agagaagccc	cgagtgggag	gggaacctgt	3480
ccaaagtcac	acggtgtgtg	ggtgacacag	ctgggggtgag	tcgaggctgg	cccctgaggc	3540
ccatgctccc	tgaacgctgg	agaccactgt	cggctagcag	cggctctcag	ggaaggcctg	3600
gtctccaccc	tcccagccta	gcctcgcgga	ccctcgctcc	ccccacatcg	gacctgtctc	3660
cctgcctgga	ccctgggctg	ccagatgcag	gaagcatcaa	accccccagc	ctcgtgggtg	3720
cggggcaggg	cgcaggcagc	acagcttaga	tgccttggtt	tgtccctctt	gtctcctggg	3780
aagagcttgc	tcccgccccag	ctctcctgcc	actggccttt	caggggtggg	ctgggcccag	3840
agtgcctttt	agtgccttct	cacggtggcc	tgatggctca	acccagtcct	aaacgggccc	3900
agtgacactg	ccgactgcac	cccagctcag	gccccactg	caccagcaat	gctagaaaac	3960
caagccaata	aaagtgattt	cttttttcat	taaaaaaaaa	aaaaa		4005

<210> 25

<211> 846

<212> DNA

<213> HOMO SAPIENS

<400> 25

caaaaatgag	cgggggtgtg	tggcccatgc	ctgtagtccc	agctgctcgg	gagactgaat	60
ctcttgagcc	tgggaagcag	aggttgcagt	gaactgagat	cgctcactg	cactccagcc	120
tgggtgacaga	gcgagattcc	atctcaaaaa	aaaaaaacag	tatgcacgta	caaatttctt	180
aacctgttat	caatgtctga	gctacataat	tatctttcta	gttggagtgt	gttttaggtg	240
tgtaccaact	gacatttcag	tttttctgtt	tgaagtccaa	tgtattagt	actctgtggc	300
tgtctctctc	acctgcccct	tgtggcctgt	ctacaattct	aaatggattt	tgaactcaat	360
gtcgtcgctt	ctgggtttct	gcatatacca	atagcattac	ctatgacttt	ttttttcctg	420
agctattttc	actgacttga	gctaattgaac	taaaactgag	ttatgtttta	tatttgtatc	480
aaatacataa	aaggaatact	gctttttcct	tttgtggctc	aaaggtagct	gcatttttaa	540
atattttgtga	aaataaaaaa	ttttgttatt	agaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	600
aaaaaaaaaa	agaaagacca	aaaaggaaga	gaaggggaaa	agaagaagag	aaacggaggg	660
acaacgggaa	acacagagag	cgagccgggtg	acgaaaaagcg	ggaaggccaa	cgcaggagaa	720
gaaagagagg	ggggcggcgt	cgctcattgt	gggagtgtcc	tcagagttaa	gcgagtgggg	780
gatgatgggc	aggagtgcata	tgcgcccctt	tgtatgaggg	ggtgcctcaa	ttgttgatgg	840
gccggg						846

<210> 26

<211> 599

<212> DNA

<213> HOMO SAPIENS

<222> 103

<400> 26

cggacggtgg	gcggacgcgt	gggaggacgc	gtgggggggtc	ttgctatgtt	taacaaccca	60
ggctgatctc	aaactcctgg	gctcaaatga	tcctccacct	tgccccctca	aaagtatggg	120
attataggca	tgagccactg	gctggcatca	ggtgccaaga	tttctgtact	gcctctaatt	180
tctgtacca	cttaaactca	ggcaggtgga	gcctacacac	tgatatttcc	ttgtggatat	240
cacacttcag	aacgtgtccg	ctagataaag	ctctcaaact	taccaaggaa	agtgatgaca	300
gcttgactcg	gccttacaca	gaaccctatg	taggtctcac	acaatagaac	aatgtacaaa	360
taagcatttt	tctttcccaa	agaagcatgt	aaagatttcc	cattcctgcc	actcaacttc	420
tccttgctgt	gacaggtggg	aagaattact	gtatatagaa	aagatgtccg	cagcgttcag	480
taaacacaga	cactaatgag	actcagaggc	tcattctgtg	tcaggtatta	taacagctta	540
aaactaaaaa	aaaaaaaaaa	aaaagggcgg	tccaagctta	ttccctttag	tgagggttaa	599

<210> 27
 <211> 603
 <212> DNA
 <213> HOMO SAPIENS

<400> 27
 gttccacggt gcttgaaatt gaaaatcaag ataaaaatgt tcacaattaa gtccttctt 60
 tttattgttc ctctagtatt ttctctcaga attgatcaag acaattcatc atttgattct 120
 ctatctccag agccaaaatc aagatttgct atgtagacg atgtaaaaat ttttagccaat 180
 ggcctccttc agttgggaca tggctctaaa gactttgtcc ataagacgaa gggccaaatt 240
 aatgacatat ttcaaaaact caacatattt gatcagctt tttatgatct atcgctgcaa 300
 accagtgaat tcaaagaaga agaaaaggaa ctgagaagaa ctacatataa actacaagtc 360
 aaaaatgaag aggtaaagaa tatgtcactt gaactcaact caaaacttga aagcctccta 420
 gaagaaaaaa ttctacttca acaaaaagtg aaatatttag aagagcaact aactaactta 480
 attcaaatc aacctgaac tccagaacac ccagaagtaa cttcacttaa agtaagtaga 540
 aaataaagag ggttcattgt tatgttttca atgtggatct tttaaaaaaa atatttctaa 600
 gcc 603

<210> 28
 <211> 879
 <212> DNA
 <213> HOMO SAPIENS

<400> 28
 gccacgcgtc cgcaaacaca aaaagaaaac aaacaaacaa aaaaacaaaa agactggctg 60
 gcggggaagg tgactcgggc ctttgctccc gagccagagc ccccaaccct gacctgatcc 120
 cctctctgct gcaggtggag ttctacttcc ttcccagta cgtgtcgcca gccgactccc 180
 cgttccgcca catcttcatg ggccgtggag accacacgct gggcgccctg ctggaccacc 240
 tgccgctgct gcgctccaac agctccggga ccccgggggc cacctcctcc actggcttcc 300
 aggagagccg tttccggcgt cagctagccc tgctcacctg gacgtgcaa ggggcagcca 360
 atgcgcttag cggggatgtc tggaacattg ataacaact ctgaggccct ggggatcctc 420
 acatccccgt cccccagtca agagctcctc tgctcctcgc ttgaatgatt cagggtcagg 480
 gaggtggctc agagtccacc tctcattgct gatcaatttc tcattacccc tacacatctc 540
 tccacggagc ccagacccca gcacagatat ccacacaccc cagccctgca gtgtagctga 600
 ccctaattgt acggtcatac tgtcgggtta tccagagata gcaccccttc aatcacagcc 660
 ccttccccct tctggggtcc tccataccta gagaccactc tgggaggttt gctaggccct 720
 gggacctggc cagctctgtt agtgggagag atcgctggca ccatagcctt atggccaaca 780
 ggtggtctgt ggtgaaaggg gcgtggagtt tcaatatcaa taaaccact gatatcaata 840
 aaaaaaaaaa aaaaaaatc tgggcgcaag aaatcgctg 879

<210> 29
 <211> 397
 <212> DNA
 <213> HOMO SAPIENS
 <222> 319, 331

<400> 29
 cctactcaac aggggggtccc aaatgcccac tgcaactagg tacagggtct gtgtgtgggt 60
 gtggaggatt ggcattggga gacatgggag gcaaagagct gggcctggcc aggcaggcc 120
 tctggcttcc aagaactcct agttccaggg gacacccagt gggggaagat ctggctgctg 180
 ggaggccac agcctagggc tggtcggcca aacagccagc tctggctcct gctcacaagt 240
 gccctatggc ttctaattga ttctcttct tctcccttg ctgcacctgc agatgcagaa 300
 ggcaggccct cccagcanc ctactggctg nccactcca tttggactgg cacattggac 360
 tggggcatca cattcctca gaacagcctg ataatg 397

<210> 30
 <211> 1740
 <212> DNA
 <213> HOMO SAPIENS
 <222> 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229,
 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244,
 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259,
 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274,

275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289,
 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304,
 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319,
 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334,
 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349,
 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364,
 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379,
 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 846, 847, 848, 849,
 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864,
 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879,
 880, 881, 882, 883, 884, 885, 886, 887, 888, 889

<400> 30

tacaaagttt	taagaaagcc	agcatctcag	aaaggccttt	caaacaagga	cacttaatta	60
gccatcttat	gtataagaaa	agaaatataa	agaacatgaa	aattttaaaaa	cagatttggc	120
agttttataa	cagtctagga	ggtggtgtta	ttttttccta	ttaagaatta	gagggcaggt	180
taggaataaa	taaaatacag	tttgaaaata	atgagnnnnn	nnnnnnnnnn	nnnnnnnnnn	240
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	300
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	360
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	420
gtggctacaa	gggaccacaa	gaattacagg	gaagttacag	ggaagcaggt	ttcatctcaa	480
tattgggaga	gatttcaaac	aatcacacct	gcctgagaag	gagtgggctg	tcactaggaa	540
tttttattcc	cagtcctgca	ggaattttgt	agaagggctt	catgtgctgg	taccaatagg	600
acaggaagat	tttaatcagc	tttactatct	atgttttttt	atggaaactg	tgtgtatgta	660
tacatacat	ttccaaaaag	aaaaattaaa	tgattataga	gattatgttt	ttcagactac	720
tcacgtatct	gcttttctta	ctccccacct	ctgctgataa	ttcctagttc	attgggtttt	780
ccccacact	ggaattacct	ggggagctta	aaaaaccctg	atgcctgggt	cccaccctca	840
gagatnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	gatatgaatg	900
cagcctgggc	actgggaaat	ttaaaaaactc	cccagataat	tactgtgaca	gccaagggtg	960
agacctccta	gtctagagct	ttgctataca	cagggtgggc	cacaagccag	gagcatcagc	1020
atcacttggc	agcgcttaga	gattcagacc	ccagaccac	tgaatctgac	cctgcatttt	1080
cactagaccc	caggtgatca	gctgcactag	actcaacctc	taaaccaaga	cctccccacc	1140
tcacagtcta	tgatccttta	gtgacctca	gctgagtcct	gtgctgaact	gtgtttgttc	1200
tccttgagca	catgcccgct	gaccagggac	agactggatg	agcaagcaac	ctgctggcta	1260
tggagaagag	ccaggctggg	taaagtgttg	ctgtgactaa	gccaggatca	aagaactgcc	1320
tggtgcttgc	actggctggc	actgagcttg	ccactctgtg	aactgtgctt	ccttcccctg	1380
catggacctg	tgccctcagtc	actattatcc	gcaggccttc	tccaagggca	gccctctcct	1440
tgtttatccc	tcttaagcct	gcgtgcagga	aggcacatta	accctgtggc	cccctgcagg	1500
caggaggggtg	ttgggtgccc	ttacctacct	tgcccttttt	ctgtgacctg	aggctgtgcc	1560
gtttatgagt	aagtgatgtg	tgtctgtgtg	tgtgtctaga	agtgtgtcac	tcacctgtgtg	1620
ttattggagg	tcgtgtaacc	ccctagcttt	gagcctggtc	tcagatgttc	ctttcccctg	1680
tctctgtcca	gccgttaacg	ccccagctct	gtaataaaag	cctatcagcc	gtgcacttta	1740

<210> 31

<211> 2394

<212> DNA

<213> HOMO SAPIENS

<400> 31

aaatgtagaa	gggaaagtcc	ttcctggtag	taatggaaaa	ccgaatggac	agagaattat	60
caatggccct	caaggacaaa	agtgggttgt	ggaccttgat	cgtgggttag	tattgaatgc	120
agaaggaagg	tacctccaag	attcacatgg	aaatcctctt	cggattaaac	taggaggaga	180
tggtcgaaac	attgtagatc	tgggaaggga	ccccgtgggtg	agtcctgacg	gcctcccact	240
ctttgggagc	gggagacatg	gcacacctct	ggccaatgcc	caggataagc	caattttgag	300
tcttgaggga	aagccgctgg	tggtgcttga	ggtcatcaaa	aaaaccaccc	atccccctac	360
cactaccatg	cagcccacca	ctactacgac	gccccctgct	accactacaa	ccccgaggcc	420
caccactgcc	accaccgcgc	gcacgaccac	caggcgctca	acaaccacag	tccgaaccac	480
tacgaggaca	accaccacca	ccacccccac	accaccact	cccatcccca	cctgtccccc	540
tgggaccttg	gaacggcacg	acgatgatgg	caacctgata	atgagctcca	atgggatccc	600
agagtgtctc	gctgaagaag	atgagttctc	aggcttggag	actgacactg	cagtacctac	660
ggaagaggcc	tacgttatat	atgatgaaga	ttatgaattt	gagacgtcaa	ggccaccaac	720
caccactgag	ccttcgacca	ctgctaccac	accgagggtg	atcccagagg	aaggcgccat	780
cagttccctt	cctgaagaag	aatttgatct	ggctggaagg	aaacgatttg	ttgctcctta	840

cgtgacgtac	ctaaataaag	acccatcagc	cccgtgctct	ctgactgatg	caactggatca	900
cttccaagtg	gacagcctcg	atgaaatcat	ccccaatgac	ctgaagaaga	gtgatctgcc	960
tccccagcat	gctccccgca	acatcacctg	gggtggccgtg	gaagggttgcc	actcatttgt	1020
cattgtggac	tgggacaaaag	ccaccccagg	agatgtggtc	acaggttact	tggtttacag	1080
tgcatectat	gaagacttca	tcaggaacaa	gtgggtccact	caagcttcat	cagtaactca	1140
cttgcccatt	gagaacctaa	agcccaacac	gagggtattat	tttaaagtgc	aagcacaaaa	1200
tcctcatggc	tacggaccta	tcagcccttc	gggtctcattt	gtcaccgaat	cagataatcc	1260
tctgcttggt	gtgaggcccc	caggcggtga	gcctatctgg	atcccattcg	ctttcaaaca	1320
tgatcccage	tacacggact	gccatggacg	gcaatatgtg	aagcgcacgt	ggtatcgaaa	1380
gttcgtggga	gttgttcttt	gtaattcact	gagggtataaa	atctacctca	gtgacaacct	1440
gaaagataca	ttctacagca	ttggagacag	ctggggaaga	ggtgaagacc	attgccaatt	1500
tgtggattca	caccttgatg	gaagaacagg	gcctcagtc	tatgtagaag	ccctccctac	1560
tattcaaggc	tactatcgcc	agtatcgta	ggagcctgtc	agggttgagg	acatcggcct	1620
cggaaacccc	tactactatg	tgggctggta	cgagtgtggg	gtctccatcc	ctggaaagtg	1680
gtaatcacag	gaccgtcatg	ctgcaagctt	gccctgccca	gccccaccaa	ctaagtcgca	1740
ctaggggctg	tgagcaaaga	cagccagcat	gctcagcccc	gctgccctag	gtgccaggaa	1800
ggtcacagat	ggacactggc	cattctgggtc	atctcagtc	ggaactcagt	cccacttctt	1860
ggcctggaca	atgaacagga	ttcagttttg	ctgttaactt	tgtctctcta	cttttttttg	1920
tttgtttgta	atagcacatc	ccagagacat	cagaaaccag	caactgattc	agtgtgattt	1980
ccagactttt	taggcatgaa	attcggacac	ttcagtattt	ccaggaatag	catatgcacg	2040
ctgttcttgc	ttcatggaat	gctacatgtg	ttctgttttt	ctcatttttg	atttctccaa	2100
aactaaactg	atttaagctt	cagggtccctt	tgtatgcagt	agaaaggaat	tattaaaaac	2160
accaccaaa	aaaataaata	tatcctactt	gaaatttact	ctatggactt	acccactgct	2220
agaataaatg	tatcaaattc	tatttgtaaa	ttctcaattt	tgatatatat	atgtatatat	2280
gcataacat	atccacactt	gtctgcaaga	atattgatta	aaattgctaa	atttgtactt	2340
gttcaccaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aagg	2394

<210> 32

<211> 499

<212> DNA

<213> HOMO SAPIENS

<400> 32

ctgggatagc	aataacctgt	gaaaatgctc	ccccggctaa	tttgtatcaa	tgattatgaa	60
caacatgcta	aatcagta	tccaaagtct	aaatatgact	attacagggtc	tggggcaaat	120
gatgaagaaa	ctttggctga	taatatgtga	gcattttcca	gatggaagct	gtatccaagg	180
atgctccgga	atgttgctga	aacagatctg	tgcacttctg	ttttaggaca	gagggtcagc	240
atgccaatat	gtgtgggggc	tacggccatg	cagcgcgatg	ctcatgtgga	cggcgagcct	300
gccactgtga	gagcctgtca	gtccctggga	acgggcatga	tgttgagttc	ctgggccacc	360
tcctcaattg	agaagtggc	ggaagctggt	cctgaggcac	ttcgttggct	gctactgtat	420
atctacaagg	accgagaagt	caccaagaag	ctagtgcggc	aggcagagaa	gatgggctac	480
aaggccatat	ttgtgacag					499

<210> 33

<211> 1774

<212> DNA

<213> HOMO SAPIENS

<222> 1679, 1681, 1684, 1685, 1686, 1691, 1692, 1693, 1705, 1706, 1707, 1708, 1710, 1714, 1716, 1731, 1740, 1744

<400> 33

ctttcctaga	caaggctgaa	agggggccaac	attatttctg	aagacttcat	tattggaatt	60
ctatgggagt	gatctcactg	agctattttg	gaatagaaat	gtggctagtt	gcctgacctc	120
ccatcaatgg	ttcacgtggc	tttcaaagg	aagggaaggc	agtgtcgtact	tttggtaaaa	180
tggggcgaag	ggtccatgcc	agcaacacaa	tactcaaa	tccagatgag	ggatcagtaa	240
atacaacgtg	cctgaaaggt	ggcccttgag	cacattcttc	cggtagacat	taacttatta	300
aattgattct	gattacaaat	ataaactttg	ccccatctc	acccagtaac	aatgcaagag	360
ttgatgtcag	tctataaaa	gaagtaggaa	ctgtccctgg	ctttcaggct	ccaacatcct	420
ccccctgtca	agatgtggca	cctcaaaact	tgtgcagtc	tcatgatctt	cctgtgtctg	480
ttggccagg	taaggaggga	aggatactta	tgtgtgtgtg	tggagtgtgg	agatgatagt	540
ggtgggtgaa	cttgaaagct	agattcagtc	ctgagggaat	gttctctctg	tctgagtcta	600
cagcatctgc	ggaatggaat	gatcactctt	ccaagggtgt	cagcagggtg	tcaaaccttt	660
catatctgaa	tgtctttgcc	cttacagata	gatggctccc	caataccaga	agtgagttcg	720

gcaaagagaa	ggccacggag	aatgacccca	ttttggagag	gggtttccct	caggcctatt	780
ggagcctcct	gccgggatga	ttctgagtgt	atcacaaggc	tatgcagaaa	aagacgctgt	840
tccttaagtg	tggcccagga	atgatgtaca	taccagggaa	agaaaggaca	gcagtcacct	900
ccgacaatgc	tccgtttctat	ggaatattga	ttaaactgcat	tttggctgga	gacacccaag	960
tgaagcaatc	ttgtattttt	aatattttaa	ggcagatgta	cgctttaaat	tggcttccat	1020
ttcttcttag	aatgttgata	tatggataag	cataactaaa	cttgtcaatt	tagagtttat	1080
ttttctatgg	atactattaa	atgtctcaaa	ttgaaatttt	agcagtctgg	aattcaagct	1140
tttgagggaa	agaaggattc	actttgtata	ctaaagaaaa	aaacagcatt	gcccaataat	1200
gtgttaactt	ctcaatctgg	aaagtgtagt	gagagctaca	taatcaatag	ctacgtaate	1260
aaacttcagca	agttcctaag	ctgtggccct	ggatcccttc	actccatact	cttcagggag	1320
gtgtcaaagg	tggtaagct	tgggaggctg	aggcaggaga	atggcgtgaa	ccgggagacg	1380
gacttgcagt	gagccgagat	ccgccactga	ctccagcctg	gggaaagagc	gagactccgt	1440
tcacaaaaaa	aaagaatcaa	aaaaaaaaag	gggagccccc	ccttggtatc	ggaagaccca	1500
gtcctgtaat	tcacacaggt	tgagttcaag	gcattaagcc	ctgtaagggc	cacttcggcc	1560
cctcagagtt	gctgttctga	tccaacggaa	gccgcttaca	aatttccctt	cggaaatttg	1620
ctccggcatt	ccctaggggc	ggatatttga	agcaaaagtc	ttttaacagc	cagtgtatnc	1680
naannncggg	nnngcccttg	cggcnnnngn	ccananattg	ctccttcttc	ncctcttctn	1740
tttnttcccc	cccgtgtcga	cagggggtgt	ggtc			1774

<210> 34

<211> 4158

<212> DNA

<213> HOMO SAPIENS

<222> 3667, 3668, 3669, 3670, 3671, 3672, 3673, 3674, 3675, 3676, 3677, 3678, 3679, 3680, 3681, 3682, 3683, 3684, 3685, 3686, 3687, 3688, 3689, 3690, 3691, 3692, 3693, 3694, 3695, 3696, 3697, 3698, 3699, 3700, 3701, 3702, 3703, 3704, 3705, 3706, 3707, 3708, 3709, 3710, 3711, 3712, 3713, 3714, 3715, 3716, 3717, 3718, 3719, 3720, 3721, 3722

<400> 34

ctcccacaac	aatttcattg	ttgttagcat	atctatttct	ccatacattg	taaaactgta	60
atccttaggt	atttctaaaa	cataaagagg	agaattaagt	cagctgcaga	acaatggggc	120
tgattcttct	gctttttctc	tggaaaatct	ttcattgctt	ttgggtggaa	tttacctaga	180
ggttacaacc	acaggatgta	gcttggtctc	ttatttgcct	ttttgggaaa	ccaattaaga	240
ttaatacagg	ataaaggaaa	aaagcaatct	attcattata	taacacagtt	gtttgtatta	300
cttgttccct	gcaaaggaaa	tctgttgaat	gcttgcat	tgaattcttt	tctaatagaa	360
caaccataaa	aggcttctta	tgggtgcagca	ggaaaaaaga	tcatttttat	agctttgcat	420
tcttaacata	gcattttaaag	agcggcatga	attagaggaa	agacatggaa	cacacaggta	480
gtcggtttga	gatcatcggc	ttaaaagtat	cctaggatgg	taatgaccca	gaagtatttc	540
cagttgtcta	gtgggtgtgg	atgcaggaat	gagaagtgtt	ttctttccat	ttcctgttgg	600
acaggtggca	atcttagcag	agccactatt	tggagttgat	aactaaagat	gcaataaacg	660
tgactatgcc	ttctggtcac	cctacgacta	tttggagttc	tccaaaacct	tgtaaagggc	720
atgtcaggca	tgcagtaaaa	gcattctacaa	cttcagctgg	gcactggcag	cataggtctc	780
atcttggacc	atacagtcac	actttataga	agagagtggg	agttctccaa	aacaatatcc	840
acaacaaagt	ctgacctcac	tctgagggag	atgggaagtg	ggaggaagaa	ggactaacca	900
gctccctgga	gtaagaggaa	tttgctttcc	ctgtctgccc	accaggggct	atatgtgcca	960
ccttttcagg	tggggccaag	gaagtgatgt	cagtgtgaca	gaaggagag	ttagacctcc	1020
agacgtcagc	ctccctccca	tggggtacat	tttcaatctg	agtgttggtg	ccttagctgt	1080
gttggtatta	gcttgattgg	tgggtccgct	ggttatgagg	tgtagggagg	cagtttttgt	1140
ttagttttta	ggactttgcc	tcttcttttg	tccttagcat	aatttctagg	cagagcatcc	1200
acgaagtccg	ttttcattgc	cagctcaaga	gcgacaatca	tttacagagt	cctatgttat	1260
gttaggtgcc	ttatgtatat	tatcccaaat	ccactgcatg	gtttaaatac	aggcactgga	1320
ataataatga	aaaaggtcat	tacagtcact	gactttctgc	aggaccttaa	acatttctct	1380
ttccacaagt	ttcccttaa	tcatgtgtca	aacctctctt	cctgacggga	atgttgtgct	1440
ataatgaatc	tgcataacgc	ttgggattct	aggaggaagg	aaggttccat	ggacatgtaa	1500
gtacagcata	ttccctcag	tcttctagga	gggcagagtg	aatcccagaa	ctggtaagat	1560
tgggaatctg	agcattgccca	ctttaatctt	agaatattta	tcattttgac	acatcctgtt	1620
tttttagagag	gaaaacaaac	acagtttctg	catttggtagt	gtaaagcata	ccttgttagg	1680
aacgtgtttt	gtaagacaca	tttgggtgtg	cattctagag	catgtcaaac	tttgacttcc	1740
aaaatatatt	tagtatgatt	gttagtggtg	acatatatca	aggctttgaa	ttactgtttt	1800
tattttaattt	tcacaagaag	cacttatttt	agccatagga	aaaccaatct	gagctacaaa	1860
tagttcttta	aaataagccc	aggttattta	gctattctag	aaagtgccga	cttctttcaa	1920
gaagcaggca	ttgtaggaca	gctgagaatt	atcacatagc	ctaaattcta	gcctggcagc	1980


```

aagagtcaca tctgagatgt ccaaaaaaaaaa aaaaaaaaaa cacctgatct acattgaaag 2040
ggggtagact aacgtatgtg agaccatttt cctatttgca gttacaagggt taaagaactt 2100
tgaaggatcat tcggctgcta agaggcatgt cgaacactct gtgtggctct ttcacagtaa 2160
accctcctaa gagcagaaga cacatggctg ttagtgctcg cgtttagatt taatttctca 2220
aataaaggcc cttggctgcg tatcatttca tccagttata aactaggggt cctgcaagca 2280
ccccattct aaggggtgaat tattgaaatc agttgctatt tgatgagtc caactggccc 2340
agcaggcagg gcatttgaag tcatggatcat caaaaagaaa tgattgtttt ttgaaaagct 2400
aaatgcttaa aatgcttcta gaggggaagtc gtggggcggtg tgctcattct ctttaaaatc 2460
aggggtgttg agtttgtttt taaacatttt tataagttca tgagaaaaaa tatataaatt 2520
ctaagaacca aactgtatt cccagaaaca tgaccctcgc tgggtcttggg tccacatatt 2580
attggactct gggggacaca aagatgcctg tgacactttg gtgttgccga gttagtcaac 2640
aattattctg ggaaaaagca gaattgaatt cttctctaga tgctctacca ggggttgcca 2700
agggccacaa agcaggctaa taaattccca caggatccag acaccaggca aaattgctct 2760
aagaagccag ttactgtcat ccctctatgg ttctagaaaa aatagtacaa aaatgacagg 2820
tcattcctatg agcgatcatg caatgaaacc ccattctctg gagaagccct tgaatcagaa 2880
ttattctttt tcttgatgtc gtcagatgca gccagtttct taattttttt aaaaactgta 2940
tgtttctgtg gtatgtatat ttgtacacct aactacctgg cacttggaat tcacagcact 3000
actcagaggc aattgaataa agagaaattt aattttaaat atcaagtcct gtcaaactt 3060
tctcaaactt ctgattttat caaagggttg ccagccaata aagtgcattc caagtataca 3120
ggggagaaaag ctgactcct acagggtcct agagttttaag taattttttt gttattaata 3180
taggtaataa tttttctaatt ttttattttt tgggttccaaa tgtaaagctc cttgtgttta 3240
cctctgttta tgctattctt gacatgttta tctaaattat gtgtgctctg tgacagggtga 3300
aatgtaaatac tgggatccat agtcaagata tcataaggac ctacttccca gcctaccttt 3360
cttctcttac ctgataatga taataactcaa aataacaaca ttcaaaggaa acacaaagaa 3420
atcctgcttt cacatctcct atttcttggg ctcttaata actactgatg gtttgttcat 3480
gaaaaaaaat ttttaaatca aaagattgta cttggccctg agttgaaaaa atttcaaaaa 3540
tcaaaagttt gtacttggcc ctgagttgaa aaaaaaaatt cacattctaa gaataaacag 3600
aaaaatgttc ttcttgggaag taaataacaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3660
aaaaaaaaann nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 3720
nntcttctat agtgtcacct aaattcaatt cactggccgt cgttttacaa cgtcgtgact 3780
gggaaaaccct tggcgattacc caacttaate gccttgacgc acatccccc ttcgcaagct 3840
ggcgtaatac tggcaaggagc ccgaaccgtt gggcccttccc aacagttgag cagcctgaat 3900
ggcgaatggg acgcgccttg tagcggcgca ttaagcgagg cgggtgtggt ggttagtaga 3960
ggtgtgcccgt aaaaaataga ataatttttt ttcaagagat gagcagaatt gagtaggaat 4020
gattacgggg aggaaaagat ctagaagata gacaatagag aggagagaaa aagagggagc 4080
aggaggctga gaggaaaaga gtagaagcgt gatatgaata tatacagaaa cagaaaaagg 4140
agagagggtg agacataa

```

<210> 35
 <211> 366
 <212> PRT
 <213> HOMO SAPIENS

```

<400> 35
Met Ala Leu Arg Phe Leu Leu Gly Phe Leu Leu Ala Gly Val Asp Leu
1      5      10      15
Gly Val Tyr Leu Met Arg Leu Glu Leu Cys Asp Pro Thr Gln Arg Leu
20     25     30
Arg Val Ala Leu Ala Gly Glu Leu Val Gly Val Gly Gly His Phe Leu
35     40     45
Phe Leu Gly Leu Ala Leu Val Ser Lys Asp Trp Arg Phe Leu Gln Arg
50     55     60
Met Ile Thr Ala Pro Cys Ile Leu Phe Leu Phe Tyr Gly Trp Pro Gly
65     70     75     80
Leu Phe Leu Glu Ser Ala Arg Trp Leu Ile Val Lys Arg Gln Ile Glu
85     90     95
Glu Ala Gln Ser Val Leu Arg Ile Leu Ala Glu Arg Asn Arg Pro His
100    105    110
Gly Gln Met Leu Gly Glu Glu Ala Gln Glu Ala Leu Gln Asp Leu Glu
115    120    125
Asn Thr Cys Pro Leu Pro Ala Thr Ser Ser Phe Ser Phe Ala Ser Leu
130    135    140

```


Leu Asn Tyr Arg Asn Ile Trp Lys Asn Leu Leu Ile Leu Gly Phe Thr
 145 150 155 160
 Asn Phe Ile Ala His Ala Ile Arg His Cys Tyr Gln Pro Val Gly Gly
 165 170 175
 Gly Gly Ser Pro Ser Asp Phe Tyr Leu Cys Ser Leu Leu Ala Ser Gly
 180 185 190
 Thr Ala Ala Leu Ala Cys Val Phe Leu Gly Val Thr Val Asp Arg Phe
 195 200 205
 Gly Arg Arg Gly Ile Leu Leu Leu Ser Met Thr Leu Thr Gly Ile Ala
 210 215 220
 Ser Leu Val Leu Leu Gly Leu Trp Asp Tyr Leu Asn Glu Ala Ala Ile
 225 230 235 240
 Thr Thr Phe Ser Val Leu Gly Leu Phe Ser Ser Gln Ala Ala Ala Ile
 245 250 255
 Leu Ser Thr Leu Leu Ala Ala Glu Val Ile Pro Thr Thr Val Arg Gly
 260 265 270
 Arg Gly Leu Gly Leu Ile Met Ala Leu Gly Ala Leu Gly Gly Leu Ser
 275 280 285
 Gly Pro Ala Gln Arg Leu His Met Gly His Gly Ala Phe Leu Gln His
 290 295 300
 Val Val Leu Ala Ala Cys Ala Leu Leu Cys Ile Leu Ser Ile Met Leu
 305 310 315 320
 Leu Pro Glu Thr Lys Arg Lys Leu Leu Pro Glu Val Leu Arg Asp Gly
 325 330 335
 Glu Leu Cys Arg Arg Pro Ser Leu Leu Arg Gln Pro Pro Pro Thr Arg
 340 345 350
 Cys Asp His Val Pro Leu Leu Ala Thr Pro Asn Pro Ala Leu
 355 360 365